arXiv:0810.2188v1 [math.CV] 13 Oct 2008

DISCRETE COMPLEX ANALYSIS ON ISORADIAL GRAPHS

DMITRY CHELKAK AND STANISLAV SMIRNOV

Abstract. We study discrete complex analysis and potential theory on a large family of planar graphs, the so-called isoradial ones. Along with discrete analogues of several classical results, we prove uniform convergence of discrete harmonic measures, Green's functions and Poisson kernels to their continuous counterparts. Among other applications, the results can be used to establish universality of the critical Ising and other lattice models.

1. Introduction

1.1. **Motivation.** This paper is concerned with discrete versions of complex analysis and potential theory in the complex plane. There are many discretizations of harmonic and holomorphic functions, which have a long history. Besides proving discrete analogues of the usual complex analysis theorems, one can ask to which extent discrete objects approximate their continuous counterparts. This can be used to give "discrete" proofs of continuous theorems (see, e.g., [L-F55] for such a proof of the Riemann mapping theorem) or to prove convergence of discrete objects to continuous ones. One of the goals of our paper is to provide tools for establishing convergence of critical 2D lattice models to conformally invariant scaling limits.

There are no "canonical" discretizations of Laplace and Cauchy-Riemann operators, the most studied ones (and perhaps the most convenient) are for the square grid. There are also definitions for other regular lattices, as well as generalizations to larger families of embedded into $\mathbb{C}$ planar graphs.

We will work with isoradial graphs (or, equivalently, rhombic lattices) where all faces can be inscribed into circles of equal radii. Rhombic lattices were introduced by R. J. Duffin [Duf68] in late sixties as (perhaps) the largest family of graphs for which the Cauchy-Riemann operator admits a nice discretization, similar to that for the square lattice. They reappeared recently as isoradial graphs in the work of Ch. Mercat [Mer01] and R. Kenyon [Ken02], as the largest family of graphs were certain 2D statistical mechanical models (notably the Ising and dimer models) preserve some integrability

---

Date: October 13, 2008.

1991 *Mathematics Subject Classification.* 39A12, 52C20, 60G50.

Key words and phrases. Discrete harmonic functions, discrete holomorphic functions, discrete potential theory, isoradial graphs, random walk.

Dept. of Math. Analysis, St. Petersburg State University. Universitetskij pr. 28, Staryj Petergof, 198504 St. Petersburg, Russia.

Section de Mathématiques, Université de Genève. 2-4 rue du Lièvre, Case postale 64, 1211 Genève 4, Suisse.

E-mail addresses: `delta4@math.spbu.ru, Stanislav.Smirnov@math.unige.ch`.

The first author was partially supported by the grants FNS 200021-107588 and NSh-2409.2008.1. The second author was partially supported by the grants FNS 200020-117596 and FNS 200020-121675.

properties. Note that isoradial graphs can be quite irregular – see e.g. Fig. 1A. It was shown by R. Kenyon and J.-M. Schlenker [KSch04] that many planar graphs admit isoradial embeddings – in fact, there are only two topological obstructions. Also isoradial graphs have a well-defined mesh size δ – the common radius of the circumscribed circles.

It is thus natural to consider this family of graphs in the context of *universality* for 2D models with (conjecturally) conformally invariant scaling limits (as the mesh tends to zero).

The primary goal of our paper is to provide a "toolbox" of discrete versions of continuous results (particularly "hard" estimates) sufficient to perform a passage to the scaling limit. Of particular interest to us is the *critical Ising model*, and this paper starts a series devoted to its *universality* (which means that the scaling limit is independent of the shape of the lattice). We plan to establish convergence of certain discrete holomorphic observables, and then deduce convergence of interfaces to Oded Schramm's SLE curves and corresponding loop ensembles. See [ChSm08], [Sm06] for the strategy of our proof and [Sm07] for the square lattice case.

Our results can also be applied to other lattice models. The uniform convergence of the discrete Poisson kernel (1.3) already implies universality for the loop-erased random walks on isoradial graphs. Namely, our paper together with [LSchW04] implies that their trajectories converge to SLE(2) curves (see Sect. 3.2, especially Remark 3.6, in [LSchW04]). There are several other fields where discrete harmonic and discrete holomorphic functions defined on isoradial graphs play essential role and hence where our results may be useful: approximation of conformal maps [Bück08]; discrete integrable systems [BMS05]; and the theory of discrete Riemann surfaces [Mer07].

Local convergence of discrete harmonic (holomorphic) functions to continuous harmonic (holomorphic) functions is a rather simple fact. Moreover, it was shown by Ch. Mercat [Mer02] that each continuous holomorphic function can be approximated by discrete ones. Thus, the discrete theory is close to the continuous theory "locally." Nevertheless, until recently almost nothing was known about the "global" convergence of the functions defined in discrete domains as the solutions of some discrete *boundary value problems* to their continuous counterparts. This setup goes back to the seminal paper by R. Courant, K. Friedrichs and H. Lewy [CFL28], where convergence is established for harmonic functions with smooth Dirichlet boundary conditions in smooth domains, discretized by the square lattice, but not much progress has occurred since. For us it is important to consider discrete domains with possibly very *rough boundaries* and to establish convergence without any regularity assumptions about them. Besides being of independent interest, this is indispensable for establishing convergence to Oded Schramm's SLEs, since the latter curves are fractal.

1.2. **Preliminary definitions.** The planar graph Γ embedded in $\mathbb{C}$ is called *isoradial* iff each face is inscribed into a circle of a common radius δ. If all circle centers are inside the corresponding faces, then one can naturally embed the dual graph Γ^* in $\mathbb{C}$ isoradially with the same δ, taking the circle centers as vertices of Γ^*. The name *rhombic lattice* is due to the fact that all quadrilateral faces of the corresponding bipartite graph Λ (having the vertex set $\Gamma \cup \Gamma^*$) are rhombi with sides of length δ (see Fig. 1A). In our paper we also require the following mild but indispensable assumption:



(A) (B)

FIGURE 1. (A) An isoradial graph Γ (black vertices, solid lines), its dual isoradial graph Γ^* (gray vertices, dashed lines), the corresponding rhombic lattice or quad-graph (vertices $\Lambda = \Gamma \cup \Gamma^*$, thin lines) and the set $\diamondsuit = \Lambda^*$ (white diamond-shaped vertices). (B) Local notations near $u \in \Gamma$.

(♠) the rhombi angles are uniformly bounded away from 0 and π (in other words, all these angles belong to $[\eta, \pi-\eta]$ for some fixed $\eta > 0$).

We will often work with rhombi half-angles, denoted by θ, for which we then have

$$0 < \mathrm{const} < \theta < \frac{\pi}{2} - \mathrm{const} < \frac{\pi}{2}.$$

Note that condition (♠) implies that for each $u_1, u_2 \in \Gamma$ the Euclidean distance $|u_2 - u_1|$ and the combinatorial distance $\delta \cdot d_\Gamma(u_1, u_2)$ (where $d_\Gamma(u_1, u_2)$ is the minimal number of vertices in the path connecting u_1 and u_2 in Γ) are comparable. Below we often use the notation const for absolute positive constants that does not depend on the mesh δ or the graph structure but, in principle, *may depend on* η.

The function $H : \Omega^\delta_\Gamma \to \mathbb{R}$ defined on some subset (discrete domain) Ω^δ_Γ of Γ is called *discrete harmonic*, if

$$\sum_{s=1}^{n} \tan\theta_s \cdot (H(u_s) - H(u)) = 0 \tag{1.1}$$

at all $u \in \Omega^\delta_\Gamma$ where the left-hand side makes sense. Here θ_s denotes the half-angles of the corresponding rhombi, see also Fig. 1B for notations. As usual, this definition is closely related to the *random walk on* Γ such that the probability to make the next step from u to u_k is proportional to $\tan\theta_k$. Namely, $\mathrm{RW}(t+1) = \mathrm{RW}(t) + \xi^{(t)}_{\mathrm{RW}(t)}$, where the increments $\xi^{(t)}$ are independent with distributions

$$\mathbf{P}(\xi_u = u_k - u) = \frac{\tan\theta_k}{\sum_{s=1}^n \tan\theta_s} \quad \text{for} \quad k = 1, .., n.$$

Under our assumption all these probabilities are uniformly bounded from 0. Note that the choice of $\tan\theta_s$ as the edge weights in (1.1) gives

$$\mathbf{E}[\operatorname{Re}\xi_u] = \mathbf{E}[\operatorname{Im}\xi_u] = 0 \quad \text{and} \quad \begin{array}{l} \mathbf{E}[(\operatorname{Re}\xi_u)^2] = \mathbf{E}[(\operatorname{Im}\xi_u)^2] = T_u, \\ \mathbf{E}[\operatorname{Re}\xi_u \operatorname{Im}\xi_u] = 0, \end{array} \tag{1.2}$$

where $T_u = \delta^2 \cdot \sum_{s=1}^n \sin 2\theta_s \big/ \sum_{s=1}^n \tan\theta_s$ (see Lemma 2.2). Our results may be directly interpreted as the convergence of the hitting probabilities for this random walk. Moreover, condition (♠) implies that quadratic variations satisfy $0 < \text{const}\,\delta^2 \leqslant T_u \leqslant \delta^2$, so after a time re-parameterization according to (1.2), this random walk converges to standard 2D Brownian motion.

1.3. **Main results.** Let $\Omega^\delta_\Gamma \subset \Gamma$ be some bounded, simply connected discrete domain and $\operatorname{Int}\Omega^\delta_\Gamma$, $\partial\Omega^\delta_\Gamma$ denote the sets of interior and boundary vertices, respectively (see Sect. 2.1 for more accurate definitions). For $u \in \operatorname{Int}\Omega^\delta_\Gamma$ and $E \subset \partial\Omega^\delta_\Gamma$ the discrete harmonic measure $\omega^\delta(u; E; \Omega^\delta_\Gamma)$ is the probability of the event that the random walk on Γ starting at u first exits Ω^δ_Γ through E. Equivalently, $\omega^\delta(\cdot; E; \Omega^\delta_\Gamma)$ is the unique discrete harmonic function in Ω^δ_Γ having boundary values 1 on E and 0 on $\partial\Omega^\delta_\Gamma \setminus E$.

We prove *uniform* (with respect to the shape Ω^δ_Γ and the structure of the underlying isoradial graph) *convergence* of the basic objects of the discrete potential theory to their continuous counterparts. Namely, we consider

- discrete harmonic measure $\omega^\delta(\cdot; a^\delta b^\delta; \Omega^\delta_\Gamma)$ of arcs $a^\delta b^\delta \subset \partial\Omega^\delta_\Gamma$ (Sect. 3.3);
- the discrete Green's function $G_{\Omega^\delta_\Gamma}^\delta(\cdot; v^\delta)$, $v^\delta \in \operatorname{Int}\Omega^\delta_\Gamma$ (Sect. 3.3);
- the discrete Poisson kernel

  $$P^\delta(\cdot; v^\delta; a^\delta; \Omega^\delta_\Gamma) = \frac{\omega^\delta(\,\cdot\,; \{a^\delta\}; \Omega^\delta_\Gamma)}{\omega^\delta(v^\delta; \{a^\delta\}; \Omega^\delta_\Gamma)}, \quad a^\delta \in \partial\Omega^\delta_\Gamma, \tag{1.3}$$

  normalized at the interior point $v^\delta \in \operatorname{Int}\Omega^\delta_\Gamma$ (Sect. 3.4);
- discrete Poisson kernel $P^\delta_{o^\delta}(\cdot; a^\delta; \Omega^\delta_\Gamma)$, $a^\delta \in \partial\Omega^\delta_\Gamma$, normalized at the boundary by the discrete analogue of the condition $\frac{\partial}{\partial n} P|_{o^\delta} = -1$ (we assume that Ω^δ_Γ has the "straight" boundary near $o^\delta \in \partial\Omega^\delta_\Gamma$, see the precise definitions in Sect. 3.5).

Moreover, we also prove uniform convergence for the *discrete gradients* of these functions (which are discrete holomorphic functions defined on subsets of $\diamondsuit = \Lambda^*$, see Sect. 2.4 and Definition 3.6 for further details).

1.4. **Organization of the paper.** We begin with the exposition of basic facts concerning discrete harmonic and discrete holomorphic functions on isoradial graphs. The larger part of Sect. 2 follows [Duf68], [Mer01], [Ken02], [Mer07] and [Bück08]. Unfortunately, none of these papers contains all the preliminaries that we need. Besides, the basic notation (sign and normalization of the Laplacian, definition of the discrete exponentials and so on) varies from source to source, so for the convenience of the reader we collected all preliminaries in the same place. Note that our notation (e.g., the additive normalization of the discrete Green's functions) is chosen to be as close in the limit to the standard continuous objects as possible. Also, we prefer to deal with functions rather than to use the language of forms or cochains [Mer07] which is more adapted for the topologically nontrivial cases.

The main part of our paper is Sect. 3, where the convergence theorems are proved. The proofs essentially use compactness arguments, so it does not give any estimate for the convergence rate. Thus, as in [Sm07], we derive the "uniform" convergence from the "pointwise" one, using the compactness of the set of bounded simply connected domains in the Carathédory topology (see Proposition 3.7). The other ingredients are the classical Arzelà-Ascoli theorem, which allows us to choose a convergent subsequence of discrete harmonic functions (see Proposition 3.1) and the weak Beurling-type estimate (Proposition 2.10) which we use in order to identify the boundary values of the limiting harmonic function. We prove C^1-convergence, but stop short of discussing the C^∞ topology since there is no straightforward definition of the second discrete derivative for functions on isoradial graphs (see Sect. 2.5). Note however that a way to overcome this difficulty was suggested in [Bück08].

Acknowledgments. We would like to thank Vincent Beffara for many helpful comments. Parts of this paper were written at the Mathematisches Forschungsinstitut Oberwolfach during the Oberwolfach-Leibniz Fellowship of the first author. The authors are grateful to the MFO for the hospitality.

2. Discrete harmonic and holomorphic functions. Basic facts

2.1. **Basic definitions. Approximation property.** Let $\Gamma = \Gamma^\delta$ be some (infinite) isoradial graph embedded into $\mathbb{C}$. Let $\Omega^\delta = F_{\Omega^\delta} \cup E_{\Omega^\delta} \cup V_{\Omega^\delta}$, where F_{Ω^δ}, E_{Ω^δ} and V_{Ω^δ} are sets of (open) faces, (open) edges and vertices of Γ^δ (see Fig. 2A). If $\Omega^\delta \subset \mathbb{C}$ is a (simply) connected domain (i.e., open subset of $\mathbb{C}$) and the skeleton $E_{\Omega^\delta} \cup V_{\Omega^\delta}$ is connected, we call $\boldsymbol{\Omega^\delta}$ the **continuous counterpart** of the **(simply) connected discrete domain** $\boldsymbol{\Omega^\delta_\Gamma}$. The latter is defined as the set of vertices $\Omega^\delta_\Gamma = \operatorname{Int}\Omega^\delta_\Gamma \cup \partial\Omega^\delta_\Gamma$, where the sets of *interior* and *boundary* vertices are given by

$$\operatorname{Int}\Omega^\delta_\Gamma := V_{\Omega^\delta} \quad \text{and} \quad \partial\Omega^\delta_\Gamma := \{(a; a_{\mathrm{int}}) : (a_{\mathrm{int}}a) \in E_{\Omega^\delta},\ a_{\mathrm{int}} \in V_{\Omega^\delta},\ a \notin V_{\Omega^\delta}\},$$

respectively. The reason for a more complicated definition of a boundary vertex (as a pair of an interior and an exterior vertices) is that it corresponds to a prime end in the continuous setting. In particular, the same vertex will play the role of several boundary vertices, when it can be approached from different directions inside the domain Ω^δ – see e.g. vertices b and c in the Fig. 2A). However, when no confusion arises we will often denote boundary vertices by a, not indicating explicitly the edge $(a_{\mathrm{int}}a)$ that ends at a.

For a domain $\Omega \subset \mathbb{C}$ we define its *discretization* Ω^δ (and, further, Ω^δ_Γ) by taking the (largest) connected component of Γ lying inside Ω as V_{Ω^δ}, and the set of all edges and faces incident to V_{Ω^δ} as E_{Ω^δ} and F_{Ω^δ}, respectively (see Fig. 2B, Fig. 3A). Note that, by definition, $\partial\Omega^\delta_\Gamma \cap \Omega = \emptyset$.

Further, we introduce the *weight* of a vertex $u \in \Gamma$ by

$$\mu^\delta_\Gamma(u) = \frac{\delta^2}{2} \sum_{u_s \sim u} \sin 2\theta_s, \tag{2.1}$$



(A) (B)

FIGURE 2. (A) Discrete domain. The interior vertices are gray, the boundary vertices are black and the outer vertices are white. Both b and c have two interior neighbors, and so we treat $b^{(1)} = (b; b^{(1)}_{\mathrm{int}})$, $b^{(2)} = (b; b^{(2)}_{\mathrm{int}})$ and $c^{(1)}$, $c^{(2)}$ as different elements of $\partial\Omega^\delta_\Gamma$. (B) Discrete half-plane $\mathbb{H}^\delta$ and discrete rectangle $R^\delta(S,T)$. The lower, upper and vertical parts of $\partial R^\delta_\Gamma(S,T)$ are denoted by $L^\delta_\Gamma(S)$, $U^\delta_\Gamma(S,T)$ and $V^\delta_\Gamma(S,T)$, respectively.

where θ_s are the half-angles of the corresponding rhombi (see Fig. 1B). An easy exercise in trigonometry shows that $\mu^\delta_\Gamma(u)$ is the area of the dual face $W(u) = w_1w_2..w_n$, $u \sim w_s \in \Gamma^*$.

Let $\phi : \overline{\Omega^\delta} \to \mathbb{C}$ be a Lipschitz (i.e., satisfying $|\phi(u_1) - \phi(u_2)| \leqslant C|u_1 - u_2|$) function and $\phi^\delta = \phi|_{\Omega^\delta_\Gamma}$ be its restriction to Ω^δ_Γ. Note that all points in the dual face $W(u)$ are δ-close to u, so approximating values of ϕ on $W(u)$ by its value at u we arrive at the inequality

$$\Bigg|\sum_{u\in\mathrm{Int}\,\Omega^\delta_\Gamma}\phi^\delta(u)\mu^\delta_\Gamma(u) + \sum_{a\in\partial\Omega^\delta_\Gamma}\phi^\delta(a)\mu^\delta_{\Omega^\delta_\Gamma}(a) - \iint_{\Omega^\delta}\phi(x+iy)dxdy\Bigg| \leqslant C\delta\cdot\mathrm{Area}(\Omega^\delta) \quad (2.2)$$

with the same constant C. Here $\mu^\delta_{\Omega^\delta_\Gamma}(a)$ is the area of $W(a)\cap\Omega^\delta$ (more accurately, the area of the connected component of $W(a)\cap\Omega^\delta$ intersecting with the edge (aa_{int})).

Definition 2.1. *Let Ω^δ_Γ be some connected discrete domain and $H : \Omega^\delta_\Gamma \to \mathbb{R}$. We define the* **discrete Laplacian** *of H at $u \in \mathrm{Int}\,\Omega^\delta_\Gamma$ as*

$$[\Delta^\delta H](u) := \frac{1}{\mu^\delta_\Gamma(u)}\sum_{u_s\sim u}\tan\theta_s\cdot[H(u_s)-H(u)].$$

We call H **discrete harmonic** *in Ω^δ_Γ iff $[\Delta^\delta H](u) = 0$ at all interior points $u \in \mathrm{Int}\,\Omega^\delta_\Gamma$.*

It is easy to see that discrete harmonic functions satisfy the *maximum principle*:

$$\max_{u\in\Omega^\delta_\Gamma} H(u) = \max_{a\in\partial\Omega^\delta_\Gamma} H(a).$$

Furthermore, *the discrete Green's formula*

$$\sum_{u\in\operatorname{Int}\Omega^\delta_\Gamma} [G\Delta^\delta H - H\Delta^\delta G](u)\mu^\delta_\Gamma(u) \;=\; \sum_{(a;a_{\mathrm{int}})\in\partial\Omega^\delta_\Gamma} \tan\theta_{aa_{\mathrm{int}}}\cdot[H(a)G(a_{\mathrm{int}}) - H(a_{\mathrm{int}})G(a)]$$

holds true for any two functions $G, H : \Omega^\delta_\Gamma \to \mathbb{R}$. Here $\theta_{aa_{\mathrm{int}}}$ denotes the half-angle of the rhombus having aa_{int} as a diagonal.

Lemma 2.2 (approximation property). *Let $\phi\in C^3$ be a smooth function defined in the disc $B(u,2\delta)\subset\mathbb{C}$ for some $u\in\Gamma$. Denote by ϕ^δ its restriction to Γ. Then*
(i) $\Delta^\delta\phi^\delta\equiv 0$ for all linear ϕ and $\Delta^\delta\phi^\delta\equiv\Delta\phi\equiv 2(a+c)$, if $\phi(x+iy)\equiv ax^2+bxy+cy^2$.
(ii)

$$\big|\,[\Delta^\delta\phi^\delta](u) - [\Delta\phi](u)\,\big| \leqslant \mathrm{const}\cdot\delta\cdot\max_{W(u)}|D^3\phi|.$$

Proof. We start by enumerating neighbors of u as $u_1,\dots,u_n$ and its neighbors on the dual lattice as $w_1,\dots,w_n$ – see Fig. 1B). Obviously, $\Delta^\delta\phi^\delta\equiv 0$, if ϕ is a constant. Since

$$\sum_{u_s\sim u}\tan\theta_s\cdot(u_s-u) = -i\sum_{u_s\sim u}(w_{s+1}-w_s) = 0,$$

one obtains $\Delta^\delta\phi^\delta\equiv 0$ for linear functions $x=\operatorname{Re}u$ and $y=\operatorname{Im}u$. Similarly,

$$\sum_{u_s\sim u}\tan\theta_s\cdot(u_s^2-u^2) = -i\sum_{u_s\sim u}(w_{s+1}-w_s)(u+u_s) = -i\sum_{u_s\sim u}(w_{s+1}^2-w_s^2) = 0,$$

so $\Delta^\delta\phi^\delta\equiv 0$ for $x^2-y^2=\operatorname{Re}u^2$ and $2xy=\operatorname{Im}u^2$. The result for x^2+y^2 follows from

$$\sum_{u_s\sim u}\tan\theta_s\cdot|u_s-u|^2 = 2\delta^2\sum_{u_s\sim u}\sin 2\theta_s = 4\mu^\delta_\Gamma(u),$$

thus proving (i). Finally, the Taylor formula implies (ii). □

2.2. Green's function. Dirichlet problem. Harnack lemma. Lipschitzness.

Definition 2.3. *Let $u_0\in\Gamma$. We call $H=G_\Gamma(\cdot;u_0):\Gamma\to\mathbb{R}$ the* **free Green function** *iff it satisfies the following:*

(i) $[\Delta^\delta H](u)=0$ for all $u\neq u_0$ and $\mu^\delta_\Gamma(u_0)\cdot[\Delta^\delta H](u_0)=1$;
(ii) $H(u)=o(|u-u_0|)$ as $|u-u_0|\to\infty$;
(iii) $H(u_0)=\frac{1}{2\pi}(\log\delta-\gamma_{\mathrm{Euler}}-\log 2)$, where γ_{Euler} is the Euler constant.

Remark 2.4. *We use the nonstandard normalization at u_0 (usually the additive constant is chosen such that $G(u_0;u_0)=0$) in order to have convergence to the standard continuous Green's function $\frac{1}{2\pi}\log|u-u_0|$ as the mesh δ goes to zero.*

Theorem 2.5 (improved Kenyon's theorem). *There exists a unique Green's function $G_\Gamma(\cdot;u_0)$. Moreover, it satisfies*

$$G_\Gamma(u;u_0) = \frac{1}{2\pi}\log|u-u_0| + O\left(\frac{\delta^2}{|u-u_0|^2}\right),\quad u\neq u_0, \tag{2.3}$$

uniformly with respect to the shape of the isoradial graph Γ and $u_0\in\Gamma$.

Proof. This asymptotic form for isoradial graphs was first obtained in [Ken02]. Some small improvements (the correct additive constant and the order of the remainder) were done in [Bück08] under the so-called quasicrystallic assumption (when the number of edge slopes in the corresponding rhombic embedding is finite). Actually, Kenyon's beautiful proof works well without any appeal to this assumption. We give the sketch of the proof (together with a slight but necessary modification) in Appendix A.1. □

Let Ω^δ_Γ be some bounded connected discrete domain. It is well known that for each $f:\partial\Omega^\delta_\Gamma\to\mathbb{R}$ there exists a unique discrete harmonic function H in Ω^δ_Γ such that $H|_{\partial\Omega^\delta_\Gamma}=f$ (e.g., H minimizes the corresponding Dirichlet energy, see [Duf68]). Clearly, H depends on f linearly, so

$$H(u)=\sum_{a\in\partial\Omega^\delta_\Gamma}\omega^\delta(u;\{a\};\Omega^\delta_\Gamma)\cdot f(a)$$

for all $u\in\Omega^\delta_\Gamma$, where $\omega^\delta(u;\cdot;\Omega^\delta_\Gamma)$ is some probabilistic measure on $\partial\Omega^\delta_\Gamma$ which is called **harmonic measure** at u. It is harmonic as a function of u and has a standard interpretation as the exit probability for the random walk on Γ (i.e. the measure of a set $A\subset\partial\Omega^\delta_\Gamma$ is the probability that the random walk started from u exits Ω^δ_Γ through A).

Definition 2.6. *Let* $u_0\in\operatorname{Int}\Omega^\delta_\Gamma$. *We call* $H=G_{\Omega^\delta_\Gamma}(\cdot;u_0)$ *the* **Green function in Ω^δ_Γ** *iff*

(i) $[\Delta^\delta H](u)=0$ *for all* $u\in\operatorname{Int}\Omega^\delta_\Gamma\setminus\{u_0\}$ *and* $\mu^\delta_\Gamma(u_0)\cdot[\Delta^\delta H](u_0)=1$;

(ii) $H=0$ *on the boundary* $\partial\Omega^\delta_\Gamma$.

Note that these two properties determine $G_{\Omega^\delta_\Gamma}(\cdot;u_0)$ uniquely. Namely, $G_{\Omega^\delta_\Gamma}=G_\Gamma-G^*_{\Omega^\delta_\Gamma}$, where

$$G^*_{\Omega^\delta_\Gamma}(\cdot;u_0)=\sum_{a\in\partial\Omega^\delta_\Gamma}\omega^\delta(\cdot;\{a\};\Omega^\delta_\Gamma)\cdot G_\Gamma(a;u_0)\qquad(2.4)$$

is the unique solution of the discrete boundary value problem

$$\Delta^\delta H^*=0\ \text{ in }\ \Omega^\delta_\Gamma,\qquad H^*=G(\cdot;u_0)\ \text{ on }\ \partial\Omega^\delta_\Gamma.$$

Applying Green's formula to $\omega^\delta(\cdot;\{a\};\Omega^\delta_\Gamma)$ and $G_{\Omega^\delta_\Gamma}(\cdot;u_0)$, one obtains

$$\omega^\delta(u_0;\{a\};\Omega^\delta_\Gamma)=-\tan\theta_{aa_{\mathrm{int}}}\cdot G_{\Omega^\delta_\Gamma}(a_{\mathrm{int}};u_0),\quad\text{where }\ a=(a;a_{\mathrm{int}})\in\partial\Omega^\delta_\Gamma.\qquad(2.5)$$

It was noted by U. Bücking [Bück08] that, since the remainder in (2.3) is of order $O(\delta^2|u-u_0|^{-2})$, one can use R. Duffin's ideas [Duf53] in order to derive the Harnack Lemma for discrete harmonic functions. Recall that we denote by $B^\delta_\Gamma(u_0,R)\subset\Gamma$ the discretization of the disc $B(u_0,R)\subset\mathbb{C}$.

Proposition 2.7 (**Harnack Lemma**)**.** *Let* $u_0\in\Gamma$ *and* $H:B^\delta_\Gamma(u_0,R)\to\mathbb{R}$ *be a nonnegative discrete harmonic function.*
(i) If $u_1\sim u_0$, *then*

$$|H(u_1)-H(u_0)|\leqslant\mathrm{const}\cdot\frac{\delta H(u_0)}{R}.$$

(ii) If $u_1, u_2 \in B^\delta_\Gamma(u_0,r) \subset \operatorname{Int} B^\delta_\Gamma(u_0,R)$, *then*

$$\exp\left[-\operatorname{const}\cdot\frac{r}{R-r}\right] \leqslant \frac{H(u_2)}{H(u_1)} \leqslant \exp\left[\operatorname{const}\cdot\frac{r}{R-r}\right].$$

Proof. In order to make our presentation complete, we recall briefly the arguments from [Bück08] and [Duf53] in Appendix A.2. □

Corollary 2.8 (Lipschitzness of discrete harmonic functions). *Let* H *be discrete harmonic in* $B^\delta_\Gamma(u_0,R)$ *and* $u_1, u_2 \in B^\delta_\Gamma(u_0,r) \subset \operatorname{Int} B^\delta_\Gamma(u_0,R)$. *Then*

$$|H(u_2)-H(u_1)| \leqslant \operatorname{const}\cdot\frac{M|u_2-u_1|}{R-r}, \quad \textit{where} \quad M = \max_{B^\delta_\Gamma(u_0,R)} |H(u)|.$$

Proof. By assumption (♠) we can find a path $u_1 = v_0v_1v_2...v_{k-1}v_k = u_2$, connecting u_1 and u_2 inside $B^\delta_\Gamma(u_0,r)$, such that $k \leqslant \operatorname{const}\cdot|u_2-u_1|\delta^{-1}$. Since $0 \leqslant H+M \leqslant 2M$, applying Harnack's inequality to $H+M$, one gets

$$|H(u_2)-H(u_1)| \leqslant \sum_{j=0}^{k-1} |H(v_{j+1})-H(v_j)| \leqslant \operatorname{const}\cdot\frac{|u_2-u_1|}{\delta}\cdot\frac{\delta M}{R-r}. \qquad \square$$

2.3. **Weak Beurling-type estimates.** The following simple fact is based on the approximation property (Lemma 2.2) for the discrete Laplacian on isoradial graphs.

Lemma 2.9. *Let* $u_0 \in \Gamma$, $r>0$ *and* $B^\delta_\Gamma(u_0,r)$ *be the discretization of the disc* $B(u_0,r)$. *Let* $a, b \in \partial B^\delta_\Gamma(u_0,r)$ *be two boundary points such that*

$$\arg(b-u_0) - \arg(a-u_0) \geqslant \tfrac{1}{4}\pi.$$

Then,

$$\omega^\delta(u; ab; B^\delta_\Gamma(0,r)) \geqslant \operatorname{const} > 0 \quad \textit{for all } u \in B^\delta_\Gamma(u_0, \tfrac{1}{2}r),$$

where ab *denotes the discrete counter clockwise arc from* a *to* b *(see Fig. 3A).*

Proof. Fix some smooth function $\phi_0 : B(0,\frac{3}{2}) \to \mathbb{R}$ such that
(ia) $\phi_0(\zeta) \leqslant 1$ near the arc $\{\zeta = e^{i\alpha}, \alpha \in [0, \frac{1}{4}\pi]\}$,
(ib) $\phi_0(\zeta) \leqslant 0$ near the complementary arc $\{\zeta = e^{i\alpha}, \alpha \in [\frac{1}{4}\pi, 2\pi]\}$;
(ii) ϕ_0 is subharmonic in $B(0,\frac{3}{2})$, moreover $[\Delta\phi_0](\zeta) \geqslant \operatorname{const} > 0$ for all $\zeta \in B(0,\frac{3}{2})$;
(iii) $\phi_0(\zeta) \geqslant \operatorname{const} > 0$ for all $\zeta \in B(0,\frac{2}{3})$.
Clearly, ϕ_0 exists, if the constants in (ii),(iii) are small enough. Let

$$\phi^\delta(u) := \phi\left(\frac{u-u_0}{a-u_0}\right) \quad \text{for} \quad u \in \Gamma.$$

Then, $\phi^\delta \leqslant 1$ on the discrete arc ab and $\phi^\delta \leqslant 0$ on the complementary arc ba.

If δ/r is small enough, then, due to (ii) and the approximation property (Lemma 2.2), ϕ^δ is discrete subharmonic in $B^\delta_\Gamma(u_0,r)$. Using the maximum principle, one obtains

$$\omega^\delta(u; ab; B^\delta_\Gamma(0,r)) \geqslant \phi^\delta(u) \geqslant \operatorname{const} > 0 \quad \text{for all } u \in B^\delta_\Gamma(0,\tfrac{1}{2}r).$$

If $\delta/r \geqslant \operatorname{const} > 0$, then the claim is trivial, since the random walk starting at u_0 can reach the discrete arc ab in a uniformly bounded number of steps. □



(A) (B)

Figure 3. (A) A discrete disc. The "black" polygonal boundary B and the "white" contour W are shown together with the correspondences $z \mapsto b(z)$, $z \in W \cap \diamondsuit$, and $z \mapsto w(z)$, $z \in B \cap \diamondsuit$. (B) The proof of the weak Beurling-type estimate (Proposition 2.10). The probability that the random walk makes a whole turn inside the annulus (and so hits the boundary $\partial\Omega^\delta$) is uniformly bounded away from 0 due to Lemma 2.9.

Let Ω^δ_Γ be some connected discrete domain, $u \in \Omega^\delta_\Gamma$ and $E \subset \partial\Omega^\delta_\Gamma$. We set

$$\mathrm{dist}_{\Omega^\delta_\Gamma}(u;E) = \inf\{R:\ u \text{ and } E \text{ are connected in } \Omega^\delta_\Gamma \cap B(u,R)\}.$$

The following Proposition is a discrete version of the classical Beurling estimate with power 1/2 replaced by some positive β.

Proposition 2.10 (weak Beurling-type estimates). *There exists an absolute constant $\beta > 0$ such that for any simply connected discrete domain Ω^δ_Γ, point $u \in \mathrm{Int}\,\Omega^\delta_\Gamma$ and some part of the boundary $E \subset \partial\Omega^\delta_\Gamma$ one has*

$$\omega^\delta(u;E;\Omega^\delta_\Gamma) \leqslant \mathrm{const}\cdot\left[\frac{\mathrm{dist}(u;\partial\Omega^\delta_\Gamma)}{\mathrm{dist}_{\Omega^\delta_\Gamma}(u;E)}\right]^\beta \quad \textit{and} \quad \omega^\delta(u;E;\Omega^\delta_\Gamma) \leqslant \mathrm{const}\cdot\left[\frac{\mathrm{diam}\,E}{\mathrm{dist}_{\Omega^\delta_\Gamma}(u;E)}\right]^\beta.$$

Above we set $\mathrm{diam}\,E := \delta$*, if E consists of one point.*

Proof. The proof is quite standard. We prove the first estimate, the second is similar. Let $d = \mathrm{dist}(u;\partial\Omega^\delta_\Gamma)$ and $r = \mathrm{dist}_{\Omega^\delta_\Gamma}(u;E)$. Without loss of generality, one may assume that $C \cdot \delta \leqslant d \leqslant \frac{1}{2}r$ for a sufficiently large constant C . The quantity $\omega^\delta(u;E;\Omega^\delta_\Gamma)$ is the probability that the random walk starting at u first hits the boundary of Ω^δ_Γ inside E. Using Lemma 2.9 (see Fig. 3B), it is easy to show that for each $d \leqslant r' \leqslant \frac{1}{2}r$ the probability to cross the annulus $B(u,2r') \setminus B(u,r')$ inside Ω^δ_Γ without touching its boundary is bounded above by some absolute constant $p < 1$ that does not depend on r' and the shape of Ω^δ_Γ. Hence,

$$\omega^\delta(u;E;\Omega^\delta_\Gamma) \leqslant p^{\log_2(r/d)-1} = p^{-1}\cdot(d/r)^{-\log_2 p},$$

so one can take the exponent $\beta = -\log_2 p > 0$. $\square$

2.4. **Factorization of the discrete Laplacian. Discrete holomorphic functions.** Let $\diamondsuit$ denotes the set of the rhombi centers. Above we discussed the theory of discrete harmonic functions defined on the isoradial graph Γ (or, in a similar manner, on its dual Γ^*). Now, following [Mer01] and [Ken02], we introduce the notion of *discrete holomorphic* functions. These are defined either on $\Lambda = \Gamma\cup\Gamma^*$, or on $\diamondsuit = \Lambda^*$. Note that, in contrast to similar Γ and Γ^*, Λ and $\diamondsuit$ have essentially different combinatorial properties.

Definition 2.11. *Let $z\in\diamondsuit$ be the center of the rhombus $v_1v_2v_3v_4$, where the vertices $v_1,v_3\in\Gamma$, $v_2,v_4\in\Gamma^*$ are enumerated in counter clockwise order. We define the (antisymmetric) weights of the edges (v_jz), (zv_j) by*

$$\mu_{v_jz} = i(v_{j+1}-v_{j-1}), \qquad \mu_{zv_j} = i(v_{j-1}-v_{j+1})$$

and the weight of $z\in\diamondsuit$ (the center of the rhombus with half-angles $\theta(z)$, $\frac{1}{2}\pi-\theta(z)$) by

$$\mu^\delta_\diamondsuit(z) = \mathrm{Area}(v_1v_2v_3v_4) = \delta^2\sin 2\theta(z).$$

For $u\in\Gamma$ (and, in the same way, for $w\in\Gamma^*$) we set $\mu^\delta_\Lambda(u) = \frac{1}{2}\mu^\delta_\Gamma(u) = \frac{1}{4}\sum_{z_s\sim u}\mu^\delta_\diamondsuit(z_s)$.

Clearly, formulas similar to (2.2) are fulfilled for ϕ defined on subsets of $\diamondsuit$ or Λ.

Definition 2.12. *(i) Let F be defined on (some part of) $\diamondsuit$. We introduce its discrete derivatives $\overline{\partial}^\delta F$ and $\partial^\delta F$ (defined at $v\in\Lambda$) as*

$$[\overline{\partial}^\delta F](v) = \frac{1}{4\mu^\delta_\Lambda(v)}\sum_{z_s\sim v}\mu_{vz_s}F(z_s) \quad \textit{and} \quad [\partial^\delta F](v) = \frac{1}{4\mu^\delta_\Lambda(v)}\sum_{z_s\sim v}\overline{\mu_{vz_s}}F(z_s).$$

We call F **discrete holomorphic at v** *iff $[\overline{\partial}^\delta F](v) = 0$.*

(ii) Let H be defined on (some part of) Λ. We define $\overline{\partial}^\delta H$ and $\partial^\delta H$ (at $z\in\diamondsuit$) as

$$\begin{aligned}[\overline{\partial}^\delta H](z) &= \frac{1}{4\mu^\delta_\diamondsuit(z)}\sum_{v_j\sim z}\mu_{zv_j}H(v_j) &= \frac{1}{2}\left[\frac{H(v_1)-H(v_3)}{\overline{v}_1-\overline{v}_3}+\frac{H(v_2)-H(v_4)}{\overline{v}_2-\overline{v}_4}\right],\\ [\partial^\delta H](z) &= \frac{1}{4\mu^\delta_\diamondsuit(z)}\sum_{v_j\sim z}\overline{\mu_{zv_j}}H(v_j) &= \frac{1}{2}\left[\frac{H(v_1)-H(v_3)}{v_1-v_3}+\frac{H(v_2)-H(v_4)}{v_2-v_4}\right].\end{aligned}$$

We call H **discrete holomorphic at z** *iff $[\overline{\partial}^\delta H](z) = 0$. We use the same notations, if H is defined on Γ (or Γ^*) only. In this case we formally set $H|_{\Gamma^*}\equiv 0$ (or $H|_\Gamma\equiv 0$).*

This definition is the natural discretization of the identities

$$\iint_P(\overline{\partial}F)(x+iy)dxdy = -\frac{i}{2}\oint_{\partial P}F(\zeta)d\zeta \quad\text{and}\quad \iint_P(\partial F)(x+iy)dxdy = \frac{i}{2}\oint_{\partial P}F(\zeta)d\overline{\zeta},$$

where P denotes the dual face $w_1w_2..w_n$ containing v or the rectangle $\widetilde{v}_1\widetilde{v}_2\widetilde{v}_3\widetilde{v}_4$, where $\widetilde{v}_j = v_j+v_{j+1}-z$ (note that the area of P is $\mu^\delta_\Gamma(v) = 2\mu^\delta_\Lambda(v)$ or $2\mu^\delta_\diamondsuit(z)$, respectively). Thus, $\overline{\partial}^\delta$ and ∂^δ have approximation properties similar to those in Lemma 2.2. Namely,

$$\begin{aligned} \left|[\overline{\partial}^\delta\phi|_\diamondsuit](v) - (\overline{\partial}\phi)(v)\right|, \ \left|[\partial^\delta\phi|_\diamondsuit](v) - (\partial\phi)(v)\right| &= O(\delta), \qquad v\in\Lambda, \\ \left|[\overline{\partial}^\delta\phi|_\Lambda](z) - (\overline{\partial}\phi)(z)\right|, \ \left|[\partial^\delta\phi|_\Lambda](z) - (\partial\phi)(z)\right| &= O(\delta^2), \qquad z\in\diamondsuit, \end{aligned}$$

for smooth functions ϕ (the additional cancellation for $\partial^\delta(\phi|_\Lambda)$ and $\overline{\partial}^\delta(\phi|_\Lambda)$ comes from the symmetry of the rectangle and, in general, does not hold for $\overline{\partial}^\delta(\phi|_\diamondsuit)$, $\partial^\delta(\phi|_\diamondsuit)$). The following factorization of the discrete Laplacian was noted in [Mer01] and [Ken02].

Proposition 2.13. *For functions H defined on subsets of Λ the following is fulfilled:*

$$[\Delta^\delta H](u) = 4[\partial^\delta\overline{\partial}^\delta H](u) = 4[\overline{\partial}^\delta\partial^\delta H](u)$$

at all points $u\in\Lambda$ where the right-hand sides make sense.

Proof. Straightforward computations show (see Fig. 1B)

$$[\overline{\partial}^\delta\partial^\delta H](u) = \frac{1}{8\mu^\delta_\Lambda(u)}\sum_{s=1}^{k}\left[\tan\theta_s\cdot[H(u_s)-H(u)] - i\cdot[H(w_{s+1})-H(w_s)]\right] = \frac{[\Delta^\delta H](u)}{4}$$

and similarly for $[\partial^\delta\overline{\partial}^\delta H](u)$. □

2.5. **Elementary properties of discrete holomorphic functions.** Let H and F be defined on some subsets of Γ (or Γ^* or Λ) and $\diamondsuit$, respectively.

1. If H is holomorphic on Λ, then H is harmonic, i.e. $H|_\Gamma$ and $H|_{\Gamma^*}$ are harmonic.
2. Conversely, in simply connected domains, H is harmonic on Γ if and only if there exists a harmonic function $\widetilde{H}$ on Γ^* such that $H+i\widetilde{H}$ (as a function on Λ) is holomorphic. $\widetilde{H}$ is defined uniquely up to an additive constant.
3. If H is harmonic on Λ, then its derivative $F=\partial^\delta H$ is holomorphic on $\diamondsuit$.
4. In simply connected domains, if F is holomorphic on $\diamondsuit$, then there exists a function $H=\int^\delta F(z)d^\delta z$ harmonic on Λ such that $\partial^\delta H = F$. Its components $H|_\Gamma$ and $H|_{\Gamma^*}$ are defined uniquely up to additive constants by
$$H(u_2)-H(u_1) = F\left(\tfrac{1}{2}(u_2+u_1)\right)\cdot(u_2-u_1), \quad u_2\sim u_1,$$
where $u_1,u_2\in\Gamma$ or $u_1,u_2\in\Gamma^*$, respectively.
5. F is holomorphic on $\diamondsuit$ if and only if both functions
$$[\mathcal{B}F](z) := \Pr\left[F(z);\overline{u_1(z)-u_2(z)}\right], \quad [\mathcal{W}F](z) := \Pr\left[F(z);\overline{w_1(z)-w_2(z)}\right]$$
are holomorphic, where $u_{1,2}(z)\in\Gamma$ and $w_{1,2}(z)\in\Gamma^*$ are the black and white neighbors of $z\in\diamondsuit$, respectively, and $\Pr[\cdot;\xi]$ denotes the orthogonal projection onto the line $\xi\mathbb{R}$ (note that $F=\mathcal{B}F+\mathcal{W}F$, since $u_1(z)-u_2(z)\perp w_1(z)-w_2(z)$).
6. If H is real and harmonic on Γ (or Γ^*) and $F=\partial^\delta H$ (recall that F is holomorphic on $\diamondsuit$), then $F=\mathcal{B}F$ (or $F=\mathcal{W}F$, respectively).
7. If H is holomorphic on Λ, then the averaged function
$$[m^\delta H](z) := \frac{1}{4}[H(u_1)+H(u_2)+H(u_3)+H(u_4)], \quad z\sim u_j\in\Lambda \qquad (2.6)$$
is holomorphic on $\diamondsuit$.

Proof. (1) Easily follows by writing $\Delta^\delta H = 4\partial^\delta\overline{\partial}^\delta H = 0$.
(2) $[\Delta^\delta H](u) = 0$ iff the sum of increments $\widetilde{H}(w_{s+1})-\widetilde{H}(w_s)$ around u is zero.
(3) Easily follows by writing $\overline{\partial}^\delta F = \frac{1}{4}\overline{\partial}^\delta\partial^\delta H = 0$.
(4) H is well-defined due to $\overline{\partial}^\delta F = 0$. By definition, $\partial^\delta H = F$, so H is harmonic.
(5) Clear, since $\mathrm{Re}[\overline{\partial}^\delta F] = \overline{\partial}^\delta[\mathcal{B}F]$ and $\mathrm{Im}[\overline{\partial}^\delta F] = \overline{\partial}^\delta[\mathcal{W}F]$.
(6) Directly follows from definitions.
(7) Straightforward calculations show $[\overline{\partial}^\delta m^\delta H](u) = 0$, since $[\overline{\partial}^\delta H](z_s) = 0$ implies

$$[m^\delta H](z_s) = \frac{H(u)}{2} + \frac{H(w_{s+1})(w_{s+1}-u) - H(w_s)(w_s-u)}{2(w_{s+1}-w_s)}\,. \qquad \square$$

Below we will also need the averaging operator m^δ (adjoint to (2.6)) for functions defined on (some part of) $\diamondsuit$:

$$[m^\delta F](v) := \frac{1}{4\mu_\Lambda^\delta(v)} \sum_{v\sim z_s\in\diamondsuit} \mu_\diamondsuit^\delta(z_s)F(z_s), \quad v \in \Lambda. \tag{2.7}$$

Unfortunately, there are several unpleasant facts that make discrete complex analysis on rhombic lattices more complicated than the standard continuous theory and even than the square lattice discretization:

- One cannot (pointwise) multiply discrete holomorphic functions: the product FG is not necessary holomorphic if both F and G are holomorphic.
- One cannot differentiate discrete holomorphic functions infinitely many times: $\partial^\delta F$ is not necessary holomorphic (on Λ) if F is holomorphic on $\diamondsuit$. Similarly, $\partial^\delta\partial^\delta H$ is not necessary holomorphic (on Λ) if H is holomorphic on Λ.
- There are no "local" discrete analogues of ∂ that map holomorphic functions on Λ (or on $\diamondsuit$) to holomorphic functions defined on the same set (Λ or $\diamondsuit$). One cannot use the natural combination of m^δ and ∂^δ since, in general, ∂^δ does not map $\mathrm{Hol}(\diamondsuit)$ into $\mathrm{Hol}(\Lambda)$ and m^δ does not map $\mathrm{Hol}(\Lambda)$ into $\mathrm{Hol}(\diamondsuit)$.

The first obstacle (multiplication) exists in all discrete theories. Concerning the second, note that in our case there is some "nonlocal" discrete differentiation (so-called dual integration, see [Duf68] and [Mer07]). Also in two particular cases the local differentiation leads to holomorphic function again: for the classical definition on the square grid (see the book by J. Lelong-Ferrand [L-F55]) and for a particular definition on the triangular lattice [DN03].

2.6. **The Cauchy kernel. The Cauchy formula. Lipschitzness.** The following asymptotic form of the discrete Cauchy kernel is due to R. Kenyon.

Theorem 2.14 (Kenyon). *Let $z_0 \in \diamondsuit$. There exists a unique function $F = K(\cdot; z_0) : \Lambda \to \mathbb{C}$ such that*

(i) $[\overline{\partial}^\delta F](z) = 0$ for all $z \neq z_0$ and $\mu_\diamondsuit^\delta(z_0) \cdot [\overline{\partial}^\delta F](z_0) = 1$;

(ii) $|F(u)| \to 0$ as $|u-z_0| \to \infty$.

Moreover, the following asymptotics holds:

$$K(u; z_0) = \frac{2}{\pi}\,\mathrm{Pr}\left[\frac{1}{u-z_0}\,;\,\overline{u_0^{(1)}-u_0^{(2)}}\right] + O\biggl(\frac{\delta}{|u-z_0|^2}\biggr), \quad u \in \Gamma;$$

$$K(w;z_0)=\frac{2}{\pi}\Pr\left[\frac{1}{w-z_0}\,;\,\overline{w_0^{(1)}-w_0^{(2)}}\right]+O\left(\frac{\delta}{|w-z_0|^2}\right),\quad w\in\Gamma^*,$$

where $u_0^{(1)},u_0^{(2)}\in\Gamma$ *and* $w_0^{(1)},w_0^{(2)}\in\Gamma^*$ *are the black and white neighbors of* z_0*, respectively.*

Proof. We give a short sketch of R. Kenyon's arguments [Ken02] (together with a slight but necessary modification) in Appendix A.1. □

Let Ω^δ_Γ be a bounded simply connected discrete domain (see Fig. 2A, 3A). Denote by $B=u_0u_1u_2..u_n$, $u_s\in\Gamma$, its closed polyline boundary, enumerated in counter clockwise order. Denote by $W=w_0w_1w_2..w_m$, $w_s\in\Gamma^*$, the closed polyline path passing throw the centers of all faces touching B (enumerated in counter clockwise order). For functions G defined on $B\cap\diamondsuit$ (or, in the same way, on $W\cap\diamondsuit$) we set

$$\oint_B^\delta G(z)d^\delta z:=\sum_{s=0}^{n-1}G\left(\tfrac{1}{2}(u_{s+1}+u_s)\right)\cdot(u_{s+1}-u_s).$$

Lemma 2.15 (Cauchy formula). *Let* $\Omega^\delta_\diamondsuit=\overline{\Omega^\delta}\cap\diamondsuit$*,* $F:\Omega^\delta_\diamondsuit\to\mathbb{C}$ *be a discrete holomorphic function (i.e.,* $(\overline{\partial}^\delta F)(v)=0$ *for all* $v\in\Omega^\delta\cap\Lambda$*) and* $z_0\in\Omega^\delta_\diamondsuit\setminus(B\cup W)$*. Then*

$$F(z_0)=\frac{1}{4i}\left[\oint_B^\delta K(w(z);z_0)F(z)d^\delta z+\oint_W^\delta K(u(z);z_0)F(z)d^\delta z\right],$$

where $z\sim w(z)\in W$*, if* $z\in B\cap\diamondsuit$*, and* $z\sim u(z)\in B$*, if* $z\in W\cap\diamondsuit$ *(see Fig. 3A).*

Proof. Discrete integration by parts and $[\overline{\partial}^\delta F](z)\equiv 0$ give

$$\begin{aligned}F(z_0)&=\frac{1}{4}\sum_{z\in\operatorname{Int}\Omega^\delta_\diamondsuit}F(z)\sum_{z\sim v\in\Lambda}\mu_{zv}K(v;z_0)=-\frac{1}{4}\sum_{v\in B\cup W}K(v;z_0)\sum_{v\sim z\in\Omega^\delta_\diamondsuit}\mu_{vz}F(z)\\&=-\frac{1}{4}\sum_{v=u(z),\,z\in W}K(v;z_0)\mu_{vz}F(z)+\frac{1}{4}\sum_{v=w(z),\,z\in B}K(v;z_0)\mu_{vz}F(z).\end{aligned}$$

It's easy to see that $\mu_{u(z)z}=i(w_{s+1}-w_s)$ for all $z=\frac{1}{2}(w_{s+1}+w_s)\in W\cap\diamondsuit$ and $\mu_{w(z)z}=-i(u_{s+1}-u_s)$ for all $z=\frac{1}{2}(u_{s+1}+u_s)\in B\cap\diamondsuit$. □

If $F=\mathcal{B}F$, i.e., F is the discrete gradient of some discrete harmonic function $H:\Omega^\delta_\Gamma\to\mathbb{R}$ (or, in a similar manner, if $F=\mathcal{W}F$), then the Cauchy formula may be nicely rewritten in the asymptotic form:

Corollary 2.16. *Let* $F=\mathcal{B}F:\Omega^\delta_\diamondsuit\to\mathbb{C}$ *be discrete holomorphic and* $z_0\in\Omega^\delta_\diamondsuit\setminus(B\cup W)$*. Then*

$$F(z_0)=\Pr\left[\frac{1}{2\pi i}\left(\oint_B^\delta\frac{F(z)d^\delta z}{z-z_0}+\oint_W^\delta\frac{F(z)d^\delta z}{z-z_0}\right)\,;\,\overline{u_0^{(1)}-u_0^{(2)}}\right]+O\left(\frac{M\delta L}{d^2}\right),$$

where $d=\operatorname{dist}(z_0,W)$*,* $M=\max_{z\in B\cup W}|F(z)|$ *and* L *is the length of* $B\cup W$*.*

Proof. We plug Kenyon's asymptotics (Theorem 2.14) of $K(v;z_0)$ into Lemma 2.15.

If $z \in B \cap \diamondsuit$, then $F(z)d^\delta z \in \mathbb{R}$, so

$$K(w(z);z_0)\cdot\frac{F(z)d^\delta z}{4i} = \Pr\biggl[\frac{F(z)d^\delta z}{2\pi i(z-z_0)}\ ;\ \overline{u_0^{(1)}-u_0^{(2)}}\,\biggr] + O\biggl(\frac{M\delta|d^\delta z|}{d^2}\biggr),$$

since $\Pr[f;\tau]\cdot\xi = \Pr[f\xi; i\tau]$ for $\xi \in i\mathbb{R}$ and $w_0^{(1)}-w_0^{(2)} \perp u_0^{(1)}-u_0^{(2)}$.

If $z \in W \cap \diamondsuit$, then $F(z)\cdot d^\delta z \in i\mathbb{R}$ and

$$K(u(z);z_0)\cdot\frac{F(z)d^\delta z}{4i} = \Pr\biggl[\frac{F(z)d^\delta z}{2\pi i(z-z_0)}\ ;\ \overline{u_0^{(1)}-u_0^{(2)}}\,\biggr] + O\biggl(\frac{M\delta|d^\delta z|}{d^2}\biggr),$$

since $\Pr[f;\tau]\cdot\xi = \Pr[f\xi;\tau]$ for $\xi \in \mathbb{R}$. Summarizing, one obtains the result. $\square$

Proposition 2.17 (Lipschitzness of discrete holomorphic functions). *Let $u_0 \in \Gamma$ and let $F = \mathcal{B}F$ be discrete holomorphic in $B^\delta_\diamondsuit(u_0,R)$. Then*

$$\bigl|F(z_s) - \Pr\bigl[2[m^\delta F](u_0);\overline{u_s-u_0}\,\bigr]\bigr| \leqslant \mathrm{const}\cdot\frac{M\delta}{R}, \quad \textit{where} \quad M = \max_{B^\delta_\diamondsuit(u_0,R)}|F(z)|,$$

for all $z_s = \frac{1}{2}(u_s+u_0) \sim u_0$. Furthermore, if $u_1, u_2 \in B^\delta_\Gamma(u_0,r) \subset \mathrm{Int}\, B^\delta_\Gamma(u_0,R)$, then

$$\bigl|[m^\delta F](u_2) - [m^\delta F](u_1)\bigr| \leqslant \mathrm{const}\cdot\frac{M|u_2-u_1|}{R-r},$$

where the averaging operator m^δ is defined by (2.7)

Proof. Let B and W be the same discrete contours as above (see Fig. 3A). Note that their lengths are bounded by $\mathrm{const}\cdot R$. Applying the asymptotic Cauchy formula (Corollary 2.16) for all $z_s \sim u_0$, one obtains

$$F(z_s) = \Pr[A;\overline{u_s-u_0}\,] + O\biggl(\frac{M\delta}{R}\biggr), \quad \text{where} \quad A = \frac{1}{2\pi i}\biggl(\oint_B^\delta \frac{F(z)d^\delta z}{z-u_0} + \oint_W^\delta \frac{F(z)d^\delta z}{z-u_0}\biggr).$$

Due to the identity

$$\frac{1}{4\mu^\delta_\Lambda(u)}\sum_{z_s\sim u}\mu^\delta_\diamondsuit(z_s)\Pr[A;\overline{u_s-u_0}\,] = \frac{1}{4\mu^\delta_\Lambda(u)}\sum_{z_s\sim u}\delta^2\sin 2\theta_s\cdot\frac{A+e^{-2i\arg(u_s-u)}\overline{A}}{2}$$

$$= \frac{A}{2} + \frac{\delta^2\overline{A}}{16i\mu^\delta_\Lambda(u)}\sum_{u_s\sim u}\bigl(e^{-2i(w_s-u_0)} - e^{-2i(w_{s+1}-u_0)}\bigr) = \frac{A}{2}$$

(see Fig. 1B), it gives

$$[m^\delta F](u_0) = \frac{A}{2} + O\biggl(\frac{M\delta}{R}\biggr).$$

In particular, $\bigl|F(z_s) - \Pr[2[m^\delta F](u_0);\overline{u_s-u_0}]\bigr| \leqslant \mathrm{const}\cdot M\delta/R$. Using similar calculations for $[m^\delta F](u_s)$ and the trivial estimate $|(z-u_s)^{-1} - (z-u_0)^{-1}| \leqslant \mathrm{const}\cdot\delta/R^2$, one obtains

$$\bigl|[m^\delta F](u_s) - [m^\delta F](u_0)\bigr| \leqslant \mathrm{const}\cdot\frac{M\delta}{R}, \quad \text{if } u_s \sim u_0.$$

Clearly, the estimate for $|[m^\delta F](u_2) - [m^\delta F](u_1)|$ follows by summation along the path (with length $\leqslant \mathrm{const}\cdot|u_2-u_1|/\delta$, which exists by condition (♠)) connecting u_1 and u_2 inside $B^\delta_\Gamma(u_0,r)$. $\square$

3. Convergence theorems

3.1. **Precompactness in the $\mathbf{C^1}$–topology.** In the continuous setup, each uniformly bounded family of harmonic functions (defined in some common domain Ω) is precompact in the C^∞–topology. Using Corollary 2.8 and Proposition 2.17, it is easy to prove the analogue of this statement for discrete harmonic functions. Let $H^\delta : \Omega^\delta_\Gamma \to \mathbb{R}$ be defined in $\Omega^\delta_\Gamma \subset \Gamma^\delta$. Then, as usual, H^δ can be thought of as defined in $\Omega^\delta \subset \mathbb{C}$ by some standard continuation (say, linear on edges and harmonic inside faces).

Proposition 3.1. *Let $H^{\delta_j} : \Omega^{\delta_j}_\Gamma \to \mathbb{R}$ be discrete harmonic functions defined in discrete domains $\Omega^{\delta_j}_\Gamma$ with $\delta_j \to 0$. Let $\Omega \subset \bigcup_{n=1}^{+\infty} \bigcap_{j=n}^{+\infty} \Omega^{\delta_j} \subset \mathbb{C}$ be some continuous domain. If H^{δ_j} are uniformly bounded on Ω, i.e.*

$$\max_{u \in \Omega^{\delta_j}_\Gamma \cap \Omega} |H^{\delta_j}(u)| \leqslant M < +\infty \quad \textit{for all} \quad j,$$

then there exists a subsequence $\delta_{j_k} \to 0$ (which we denote by δ_k for shortness) and two functions $h : \Omega \to \mathbb{R}$, $f : \Omega \to \mathbb{C}$ such that (we denote by "$\rightrightarrows$" uniform convergence)

$$H^{\delta_k} \rightrightarrows h \quad \textit{uniformly on compact subsets } K \subset \Omega$$

and

$$\frac{H^{\delta_k}(u_2^k) - H^{\delta_k}(u_1^k)}{|u_2^k - u_1^k|} \rightrightarrows \operatorname{Re}\left[f(u) \cdot \frac{u_2^k - u_1^k}{|u_2^k - u_1^k|} \right], \tag{3.1}$$

if $u_1^k, u_2^k \in \Gamma^{\delta_k}$, $u_2^k \sim u_1^k$ and $u_1^k, u_2^k \to u \in K \subset \Omega$. Moreover, the limit function h is harmonic in Ω, $|h| \leqslant M$ and $f = h'_x - ih'_y = 2\partial h$ is analytic in Ω.

Remark 3.2. *In other words, the discrete gradients of H^δ defined by the left-hand side of (3.1) converge to ∇h. Looking at the edge $(u_1 u_2)$ one sees only the discrete directional derivative of H^δ along $\tau = (u_2 – u_1)/|u_2 – u_1|$ which converges to $\langle \nabla h(u), \tau \rangle = 2\partial h(u) \cdot \tau$.*

Proof. Due to the uniform Lipschitzness of bounded discrete harmonic functions (see Corollary 2.8) and the Arzelà-Ascoli Theorem, the sequence $\{H^{\delta_j}\}$ is precompact in the uniform topology on any compact subset $K \subset \Omega$. Moreover, their discrete derivatives (defined for $z \in \Omega^{\delta_j}_\diamondsuit$)

$$F^{\delta_j}(z) = \frac{H^{\delta_j}(u_2(z)) - H^{\delta_j}(u_1(z))}{u_2(z) - u_1(z)}, \qquad z = \tfrac{1}{2}(u_1(z) + u_2(z)), \quad z \sim u_{1,2}(z) \in \Gamma^{\delta_j},$$

are discrete holomorphic and bounded on any compact subset $K \subset \Omega$. Then, due to Proposition 2.17 (ii) and the Arzelà-Ascoli Theorem, the sequence of averaged functions $m^{\delta_j} F^{\delta_j}$ (defined on $\Omega^{\delta_j}_\Gamma$ by (2.7)) is precompact in the uniform topology on any compact subset of Ω. Thus, for some subsequence $\delta_k \to 0$, one has

$$H^{\delta_k} \rightrightarrows h \qquad \text{and} \qquad 2m^{\delta_k} F^{\delta_k} \rightrightarrows f$$

uniformly on compact subsets of Ω. Moreover, due to Proposition 2.17 (i), it gives

$$\left| F^{\delta_k}(z) - \Pr\left[f(z); \overline{u_2(z) - u_1(z)} \right] \right| \rightrightarrows 0 \quad \text{uniformly on compact subsets of } \Omega.$$

It's easy to see that h is harmonic. Indeed, let $\phi : \Omega \to \mathbb{R}$ be an arbitrary $C_0^\infty(\Omega)$ test function (i.e., $\phi \in C^\infty$ and $\operatorname{supp} \phi \subset \Omega$). Denote by h^δ, ϕ^δ and $(\Delta\phi)^\delta$ the restrictions of

h, ϕ and $\Delta\phi$ onto the lattice Γ^δ. The approximation properties ((2.2) and Lemma 2.2) and discrete integration by parts give

$$\langle h, \Delta\phi\rangle_\Omega \;=\; \lim_{\delta=\delta_k\to 0}\sum_{u\in\Omega_\Gamma^\delta} h^\delta(u)(\Delta\phi)^\delta(u)\mu_\Gamma^\delta(u) \;=\; \lim_{\delta=\delta_k\to 0}\sum_{u\in\Omega_\Gamma^\delta} h^\delta(u)[\Delta^\delta\phi^\delta](u)\mu_\Gamma^\delta(u)$$

$$=\lim_{\delta=\delta_k\to 0}\sum_{u\in\Omega_\Gamma^\delta} H^\delta(u)[\Delta^\delta\phi^\delta](u)\mu_\Gamma^\delta(u) \;=\; \lim_{\delta=\delta_k\to 0}\sum_{u\in\Omega_\Gamma^\delta}[\Delta^\delta H^\delta](u)\phi^\delta(u)\mu_\Gamma^\delta(u) \;=\; 0.$$

Furthermore, for any path $[u_0u_n]^\delta = u_0u_1..u_n$, $u_{s+1}\sim u_s$, $u_s\in\Gamma^\delta$, one has

$$H(u_n)-H(u_0)=\int_{[u_0u_n]^\delta}^{\delta} F^\delta(z)d^\delta z=\sum_{s=0}^{n-1} F^\delta\big(\tfrac{1}{2}(u_{s+1}+u_s)\big)\cdot(u_{s+1}-u_s).$$

Taking appropriate discrete approximations of segments $[uv]\subset\Omega$ (recall that rhombi angles are bounded from 0 and π, so one may find polyline approximations with uniformly bounded lengths) and passing to the limit as $\delta=\delta_k\to 0$, one obtains

$$h(v)-h(u)=\int_{[uv]}\mathrm{Re}[f(z)dz]=\mathrm{Re}\left[\int_{[uv]} f(z)dz\right]\quad\text{for all segments } [uv]\subset\Omega.$$

It gives $\alpha h'_x(u)+\beta h'_y(u)=\mathrm{Re}[(\alpha+i\beta)f(u)]$ for all $u\in\Omega$ and $\alpha,\beta\in\mathbb{R}$, so $f=2\partial h$. $\square$

3.2. **Carathéodory topology and uniform $\mathbf{C^1}$-convergence.**

Below we need some standard concepts of geometric function theory (see [Pom92], Chapters 1,2).

Let Ω be a simply connected domain. A *crosscut* C of Ω is an open Jordan arc in Ω such that $\overline{C}=C\cup\{a,b\}$ with $a,b\in\partial\Omega$. A *prime end* of Ω is an equivalence class of sequences (*null-chains*) (C_n) such that $\overline{C}_n\cap\overline{C}_{n+1}=\emptyset$, C_n separates C_0 from C_{n+1} and $\operatorname{diam} C_n\to 0$ as $n\to\infty$ (two null chains (C_n), $(\widetilde{C})_n$ are equivalent iff for all sufficiently large m there exists n such that C_m separates $\widetilde{C}_0$ from $\widetilde{C}_n$ and $\widetilde{C}_m$ separates C_0 from C_n).

Let $P(\Omega)$ denote the set of all prime ends of Ω and let $\phi:\Omega\to\mathbb{D}$ be a conformal map. Then (see Theorem 2.15 in [Pom92]) ϕ induces the natural bijection between $P(\Omega)$ and the unit circle $\mathbb{T}=\partial\mathbb{D}$.

Let $u_0\in\mathbb{C}$ be given and $\Omega_n,\Omega\subset\mathbb{C}$, be simply connected domains $\neq\mathbb{C}$ with $u_0\in\Omega_n,\Omega$. We say that $\Omega_n\to\Omega$ as $n\to\infty$ *in the sense of kernel convergence with respect to* u_0 iff

(i) some neighborhood of every $u\in\Omega$ lies in Ω_n for large enough n;

(ii) for every $a\in\partial\Omega$ there exist $a_n\in\partial\Omega_n$ such that $a_n\to a$ as $n\to\infty$. +

Let $\phi_k:\Omega_k\to\mathbb{D}$, $\phi:\Omega\to\mathbb{D}$ be the Riemann uniformization maps normalized at u_0 (i.e., $\phi(u_0)=0$ and $\phi'(u_0)>0$). Then (see Theorem 1.8 [Pom92])

$$\Omega_k\to\Omega \text{ w.r.t. } u_0 \qquad\Leftrightarrow\qquad \phi_k^{-1}\rightrightarrows\phi^{-1}\text{ uniformly on compact subsets of }\mathbb{D}.$$

It is easy to see that (a) $\phi_k\rightrightarrows\phi$ as $k\to\infty$ uniformly on compact subsets $K\subset\Omega$; (b) for all $0<r<R$ the set of simply connected domains $\Omega: B(u_0,r)\subset\Omega\subset B(0,R)$ is compact in the topology of kernel convergence w.r.t. u_0.

Definition 3.3. *Let $\Omega=(\Omega;v,..;a,b,..)$ be a simply connected bounded domain with several (possibly none) marked interior points $v,..\in\operatorname{Int}\Omega$ and prime ends (boundary* *points)* $a, b, .. \in P(\Omega)$ *(we admit coincident marked points, say,* $a=b$*) and let* $u \in \Omega$*. We write*

$$(\Omega_k; u_k) = (\Omega_k; u_k, v_k, ..; a_k, b_k, ..) \xrightarrow{\text{Cara}} (\Omega; u) = (\Omega; u, v, ..; a, b, ..) \quad \text{as } k \to \infty,$$

iff the domains Ω_k *are uniformly bounded,* $u_k \to u$, $\Omega_k \to \Omega$ *in the sense of kernel convergence w.r.t.* u *and* $\phi_k(v_k) \to \phi(v), ..,\ \phi_k(a_k) \to \phi(a), ..,$ *where* $\phi_k : \Omega_k \to \mathbb{D}$, $\phi : \Omega \to \mathbb{D}$ *are the Riemann uniformization maps normalized at* u*.*

Remark 3.4. *Since* $v \in \Omega$*, one has* $|\phi(v)| < 1$*. Thus,* $\phi_k(v_k) \to \phi(v)$ *implies* $v_k \to v$*. Moreover, one can equivalently use the point* v *instead of* u *in the definition given above.*

Definition 3.5. *Let* Ω *be a simply connected bounded domain,* $u, v, .. \in \Omega$ *and* $r > 0$*. We say that the inner points* $u, v, ..$ *are* **jointly r-inside Ω** *iff* $B(u, r), B(v, r), .. \subset \Omega$ *and there are paths* $L_{uv}, ..$ *connecting these points* r*-inside* Ω *(i.e.,* $\mathrm{dist}(L_{uv}, \partial\Omega), .. \geqslant r$*). In other words,* $u, v, ..$ *belong to the same connected component of the* r*-interior of* Ω*.*

Note that for each $0 < r < R$ there exists some $C(r, R)$ such that, if $\Omega \subset B(0, R)$ and $u, v, ..$ are jointly r-inside Ω, then

$$|\phi(v)|, .. \leqslant C(r, R) < 1, \tag{3.2}$$

where $\phi : \Omega \to \mathbb{D}$ is the Riemann uniformization map normalized at u. Indeed, considering the standard plane metric, one concludes that the extremal distance between L_{uv} and $\partial\Omega$ in $\Omega \setminus L_{uv}$ is no greater than $\pi R^2/r$. Thus, the conformal modulus of the annulus $\mathbb{D} \setminus \phi(L_{uv})$ is bounded below by some $\mathrm{const}(r, R) > 0$. Since $\phi(u) = 0$, (3.2) holds true for some $C(r, R) < 1$.

Suppose that a harmonic function

$$h(\,\cdot\,; \Omega) = h(\,\cdot\,, v, ..; a, b, ..; \Omega) : \Omega \to \mathbb{R}$$

is defined for each $\Omega = (\Omega; v, ..; a, b, ..)$. Similarly, let $\Omega^\delta_\Gamma = (\Omega^\delta_\Gamma; v^\delta, ..; a^\delta, b^\delta, ..)$ denote a simply connected bounded discrete domain with several marked vertices $v^\delta, .. \in \mathrm{Int}\,\Omega^\delta_\Gamma$ and $a^\delta, b^\delta, .. \in \partial\Omega^\delta_\Gamma$ and

$$H^\delta(\,\cdot\,; \Omega^\delta_\Gamma) = H^\delta(\,\cdot\,, v^\delta, ..; a^\delta, b^\delta, ..; \Omega^\delta_\Gamma) : \Omega^\delta_\Gamma \to \mathbb{R}$$

be some discrete harmonic function in Ω^δ_Γ. Recall that Ω^δ denotes the corresponding *continuous* polygonal domain.

Definition 3.6. *We say that the functions* $\mathbf{H}^\delta$ **are uniformly $\mathbf{C^1}$–close to h inside $\mathbf{\Omega}^\delta$***, iff for all* $0 < r < R$ *there exists* $\varepsilon(\delta) = \varepsilon(\delta, r, R) \to 0$ *as* $\delta \to 0$ *such that for all discrete domains* $\Omega^\delta_\Gamma = (\Omega^\delta_\Gamma; v^\delta, ..; a^\delta, b^\delta, ..)$ *and* $u^\delta \in \mathrm{Int}\,\Omega^\delta_\Gamma$ *the following holds true:*

If $\Omega^\delta \subset B(0, R)$ *and* $u^\delta, v^\delta, ..$ *are jointly* r*-inside* Ω^δ*, then*

$$\big|H^\delta(u^\delta, v^\delta, ..; a^\delta, b^\delta, ..; \Omega^\delta_\Gamma) \;-\; h(u^\delta, v^\delta, ..; a^\delta, b^\delta, ..; \Omega^\delta)\big| \leqslant \varepsilon(\delta) \tag{3.3}$$

and, for all $u^\delta \sim u^\delta_1 \in \Omega^\delta_\Gamma$*,*

$$\left|\frac{H^\delta(u^\delta_1; \Omega^\delta_\Gamma) - H^\delta(u^\delta; \Omega^\delta_\Gamma)}{|u^\delta_1 - u^\delta|} \;-\; \mathrm{Re}\Big[2\partial h(u^\delta; \Omega^\delta) \cdot \frac{u^\delta_1 - u^\delta}{|u^\delta_1 - u^\delta|}\Big]\right| \leqslant \varepsilon(\delta), \tag{3.4}$$

where $2\partial h = h'_x - ih'_y$*.*

Proposition 3.7. *Let (a) h be Carathéodory-stable, i.e.,*

$$h(u_k;\Omega_k) \to h(u;\Omega), \quad \text{if} \quad (\Omega_k;u_k) \overset{\text{Cara}}{\longrightarrow} (\Omega;u) \quad \text{as} \quad k\to\infty; \tag{3.5}$$

and (b) $H^\delta \to h$ pointwise as $\delta\to 0$, i.e.,

$$H^\delta(u^\delta;\Omega^\delta_\Gamma) \to h(u;\Omega), \quad \text{if} \quad (\Omega^\delta;u^\delta) \overset{\text{Cara}}{\longrightarrow} (\Omega;u) \quad \text{as} \quad \delta\to 0. \tag{3.6}$$

Then the functions H^δ are uniformly C^1–close to h inside Ω^δ (see Definition 3.6).

Proof. Assume (3.3) does not hold true, i.e.,

$$\left|H^\delta(u^\delta;\Omega^\delta_\Gamma) - h(u^\delta;\Omega^\delta)\right| \geqslant \varepsilon_0 > 0$$

for some sequence $(\Omega^\delta_\Gamma;u^\delta)$, $\delta=\delta_j\to 0$, such that $B(u^\delta,r)\subset\Omega^\delta\subset B(0,R)$. Taking a subsequence, one may assume that $u^\delta\to u$ for some $u\in B(0,R)$. The set of all simply connected domains $\Omega : B(u,\frac{1}{2}r)\subset\Omega\subset B(0,R)$ is compact in the Carathéodory topology. Thus, taking a subsequence again, one may assume that

$$(\Omega^\delta;u^\delta,v^\delta,..;a^\delta,b^\delta,..) \overset{\text{Cara}}{\longrightarrow} (\Omega;u,v,..;a,b,..) \quad \text{as} \quad \delta=\delta_k\to 0$$

(note that the marked points $v^\delta,..$ cannot reach the boundary due to (3.2)). Then, (a) the Carathéodory-stability of h and (b) the pointwise convergence $H^\delta\to h$ give a contradiction. Indeed, both

$$\text{(a) } h(u^\delta;\Omega^\delta)\to h(u;\Omega) \quad \text{and} \quad \text{(b) } H^\delta(u^\delta;\Omega^\delta_\Gamma)\to h(u;\Omega) \quad \text{as } \delta=\delta_k\to 0.$$

In view of Proposition 3.1, the proof for discrete gradients goes by the same way. Assume (3.4) does not hold for some sequence of discrete domains. As above, one may take a subsequence $\delta=\delta_k$ such that $(\Omega^\delta;u^\delta)\overset{\text{Cara}}{\longrightarrow}(\Omega,u)$. The Carathéodory-stability implies

$$h(\,\cdot\,;\Omega^\delta) \rightrightarrows h(\,\cdot\,;\Omega) \quad \text{uniformly on} \quad \overline{B(u,\tfrac{1}{2}r)} \quad \text{as} \quad \delta=\delta_k\to 0.$$

Indeed, $(\Omega^\delta;\widetilde{u})\overset{\text{Cara}}{\longrightarrow}(\Omega;\widetilde{u})$ for all $\widetilde{u}\in\overline{B(u,\frac{1}{2}r)}$. If $|h(\widetilde{u}^\delta;\Omega^\delta)-h(\widetilde{u}^\delta;\Omega)|\geqslant\varepsilon_0>0$ for some $\widetilde{u}^\delta$ and all $\delta=\delta_k$, then, taking a subsequence $\delta=\delta_m$ such that $\widetilde{u}^\delta\to\widetilde{u}\in\overline{B(u,\frac{1}{2}r)}$, one obtains a contradiction with $h(\widetilde{u}^\delta;\Omega^\delta)\to h(\widetilde{u};\Omega)$ and $h(\widetilde{u}^\delta;\Omega)\to h(\widetilde{u};\Omega)$.

The uniform estimate $|H^\delta(\,\cdot\,;\Omega^\delta_\Gamma)-h(\,\cdot\,;\Omega^\delta)|\leqslant\varepsilon(\delta)\to 0$ (see above) gives

$$H^\delta(\,\cdot\,;\Omega^\delta_\Gamma) \rightrightarrows h(\,\cdot\,;\Omega) \quad \text{uniformly on} \quad \overline{B(u,\tfrac{1}{2}r)} \quad \text{as} \quad \delta=\delta_k\to 0.$$

In particular, the functions $H^{\delta_k}(\cdot;\Omega^{\delta_k}_\Gamma)$ are uniformly bounded in $\overline{B(u,\frac{1}{2}r)}$. Thus, using Proposition 3.1, one can find a subsequence $\delta=\delta_m$ such that the discrete derivatives of H^{δ_m} converge (as defined by (3.4)) to $f=2\partial h(\cdot;\Omega)$ which gives a contradiction. $\square$

3.3. **Convergence for the harmonic measure of arcs and the Green's function.**

Theorem 3.8. *The discrete harmonic measures $\omega^\delta(\cdot;b^\delta a^\delta;\Omega^\delta_\Gamma)$ are uniformly C^1–close to the continuous harmonic measures $\omega(\cdot;a^\delta b^\delta;\Omega^\delta)$ inside Ω^δ (see Definition 3.6).*

Proof. By conformal invariance, the continuous harmonic measure is Carathéodory stable, so the first assumption in Proposition 3.7 holds true. Thus, it is sufficient to prove pointwise convergence (3.6). Let $(\Omega^\delta;u^\delta;a^\delta,b^\delta)\overset{\text{Cara}}{\longrightarrow}(\Omega;u;a,b)$. Obviously, the functions $\omega^\delta(\cdot;b^\delta a^\delta;\Omega^\delta_\Gamma)$ are uniformly bounded in Ω. Due to Proposition 3.1, one can find a subsequence $\delta_k\to 0$ such that

$$\omega^{\delta_k}(\cdot;b^{\delta_k}a^{\delta_k};\Omega^{\delta_k}_\Gamma)\rightrightarrows H$$

uniformly on compact subsets of Ω, where $H:\Omega\to\mathbb{R}$ is some harmonic function. It is sufficient to prove that $H(u)=\omega(u;ba;\Omega)$ for each subsequential limit.

Let $z^\delta\in\operatorname{Int}\Omega^\delta_\Gamma$. The weak Beurling-type estimate (see Proposition 2.10) gives

$$0\leqslant\omega^\delta(z^\delta;b^\delta a^\delta;\Omega^\delta_\Gamma)\leqslant\mathrm{const}\cdot\left[\frac{\operatorname{dist}(z^\delta;\partial\Omega^\delta_\Gamma)}{\operatorname{dist}_{\Omega^\delta_\Gamma}(z^\delta;b^\delta a^\delta)}\right]^\beta$$

uniformly as $\delta\to 0$. Passing to the limit as $\delta=\delta_k\to 0$, one obtains

$$0\leqslant H(z)\leqslant\mathrm{const}\cdot\left[\frac{\operatorname{dist}(z;\partial\Omega)}{\operatorname{dist}_\Omega(z;ba)}\right]^\beta\quad\text{for all } z\in\operatorname{Int}\Omega.$$

Therefore, $H\equiv 0$ on the arc $ab\subset\partial\Omega$. Similar arguments give $H\equiv 1$ on the arc $ba\subset\partial\Omega$. Hence, $H=\omega(\cdot;ba;\Omega)$ and, in particular, $H(u)=\omega(u;ba;\Omega)$. $\square$

Let Ω^δ_Γ be some bounded simply connected discrete domain. Recall that the discrete Green's function $G_{\Omega^\delta_\Gamma}(\,\cdot\,;v^\delta)$, $v^\delta\in\operatorname{Int}\Omega^\delta_\Gamma$, is given by

$$G_{\Omega^\delta_\Gamma}(\,\cdot\,;v^\delta)=G_\Gamma(\,\cdot\,;v^\delta)-G^*_{\Omega^\delta_\Gamma}(\,\cdot\,;v^\delta):\Omega^\delta_\Gamma\to\mathbb{R}$$

(see (2.4)). Theorem 2.5 claims uniform C^1–convergence of the free Green's function G_Γ to its continuous counterpart

$$G_{\mathbb{C}}(u;v)=\frac{1}{2\pi}\log|u-v|$$

with an error of order $O(\delta^2|u-v|^{-2})$ for the functions and so $O(\delta|u-v|^{-2})$ for their gradients. Let $G^*_\Omega(\cdot;v)=G_{\mathbb{C}}(\cdot;v)-G_\Omega(\cdot;v):\Omega\to\mathbb{R}$ be the solution of the Dirichlet problem

$$\Delta G^*_\Omega(\cdot;v)=0\quad\text{in }\ \Omega\qquad\text{and}\qquad G^*_\Omega(\cdot;v)=G_{\mathbb{C}}(\cdot;v)\quad\text{on }\ \partial\Omega.$$

Theorem 3.9. *The discrete harmonic functions $G^*_{\Omega^\delta_\Gamma}(\cdot;v^\delta)$ are uniformly C^1–close to their continuous counterparts $G^*_{\Omega^\delta}(\cdot;v^\delta)$ inside Ω^δ (see Definition 3.6).*

Proof. We start with the proof of (3.6). Let $(\Omega^\delta;u^\delta,v^\delta)\xrightarrow{\mathrm{Cara}}(\Omega;u,v)$ as $\delta=\delta_j\to 0$. We need to check that $G^*_{\Omega^\delta_\Gamma}(u^\delta;v^\delta)\to G^*_\Omega(u;v)$. Recall that $u^\delta\to u$, $v^\delta\to v$ and $B(u,r),B(v,r)\subset\Omega^\delta$ for some $r>0$, if δ is small enough. Let

$$g(z)=G_{\mathbb{C}}(z;v)\quad\text{for}\ \ z:|z-v|\geqslant\tfrac12 r.$$

Denote by $\widetilde{G}^*(\cdot;\Omega^\delta_\Gamma)$ the solution of the discrete Dirichlet problem

$$\Delta^\delta\widetilde{G}^*=0\quad\text{in }\ \Omega^\delta_\Gamma\qquad\text{and}\qquad\widetilde{G}^*=g\quad\text{on }\ \partial\Omega^\delta_\Gamma.$$

Due to the maximum principle, since $v^\delta\to v$ and $|G_\Gamma(\cdot;v^\delta)-G_{\mathbb{C}}(\cdot;v^\delta)|\to 0$ as $\delta\to 0$ uniformly outside $B(v,\frac12 r)$, it is sufficient to prove that

$$\widetilde{G}^*(u^\delta;\Omega^\delta_\Gamma)\to G^*_\Omega(u;v)\quad\text{as}\ \ \delta=\delta_j\to 0.$$

Note that the functions $\widetilde{G}^*(\cdot;\Omega^\delta_\Gamma)$ are uniformly bounded. It follows from Proposition 3.1 that one can find some subsequence $\delta_k\to 0$ such that

$$\widetilde{G}^*(\cdot;\Omega^{\delta_k}_\Gamma)\rightrightarrows H$$

uniformly on compact subsets of Ω, where $H:\Omega\to\mathbb{R}$ is some harmonic function. It is sufficient to prove that $H(u)=G^*_\Omega(u;v)$ for each subsequential limit H.

Let $z^\delta\in\Omega^\delta_\Gamma\cap\operatorname{Int}\Omega$, $|z^\delta-v|\geqslant r$ and $\operatorname{dist}(z^\delta;\partial\Omega)=d$. For each $d<\rho<R$ one has

$$\begin{aligned}\left|\widetilde{G}^*(z^\delta;\Omega^\delta_\Gamma)-g(z^\delta)\right| \;&\leqslant\; \max\left\{|g(w^\delta)-g(z^\delta)|,\ w^\delta\in\partial\Omega^\delta_\Gamma\cap B(z^\delta,\rho)\right\}\\ &+\;\max\left\{|g(w^\delta)-g(z^\delta)|,\ w^\delta\in\partial\Omega^\delta_\Gamma\right\}\cdot\omega^\delta(z^\delta;\partial\Omega^\delta_\Gamma\setminus B(z^\delta,\rho);\Omega^\delta_\Gamma).\end{aligned}$$

Since Ω^δ_Γ are uniformly bounded by our assumption (see Definition 3.6), the function g is uniformly bounded and Lipschitz in $\Omega^\delta\setminus B(v,\frac{1}{2}r)$. Moreover, $\operatorname{dist}(z^\delta;\partial\Omega^\delta_\Gamma)\leqslant 2d$, if δ is small enough. Applying Proposition 2.10, one obtains

$$\left|\widetilde{G}^*(z^\delta;\Omega^\delta_\Gamma)-g(z^\delta)\right|=O(\rho+(2d/\rho)^\beta)\quad\text{uniformly}\ \ \text{as}\ \ \delta=\delta_k\to 0.$$

Setting $\rho=d^{\frac{\beta}{\beta+1}}$ and passing to the limit as $\delta=\delta_k\to 0$, one obtains

$$|H(z)-g(z)|=O\left((\operatorname{dist}(z,\partial\Omega))^{\frac{\beta}{\beta+1}}\right)\quad\text{for all}\ \ z\in\Omega\setminus B(v,\tfrac{1}{2}r).$$

Thus, $H=g=G^*_\Omega(\cdot;v)$ on the boundary $\partial\Omega$. Since both H and $G^*_\Omega(\cdot;v)$ are harmonic in Ω, it gives $H(u)=G^*_\Omega(u;v)$.

Repeating the arguments given above in the continuous setup, it's easy to derive the Carathéodory stability of the function $G^*_\Omega(\cdot;v)$. Thus, both assumptions (3.5) and (3.6) in Proposition 3.7 hold true. □

3.4. Convergence for the Poisson kernel normalized at an interior point.

Let Ω^δ_Γ be some simply connected discrete domain, $a^\delta\in\partial\Omega^\delta_\Gamma$ and $v^\delta\in\operatorname{Int}\Omega^\delta_\Gamma$. We call

$$P^\delta=P^\delta(\,\cdot\,;v^\delta;a^\delta;\Omega^\delta_\Gamma):\Omega^\delta_\Gamma\to\mathbb{R}$$

the *discrete Poisson kernel normalized at* v^δ if

$$\Delta^\delta P^\delta=0\ \text{ in }\ \Omega^\delta_\Gamma,\qquad P^\delta\big|_{\partial\Omega^\delta_\Gamma\setminus\{a\}}=0\qquad\text{and}\qquad P^\delta(v^\delta)=1$$

Note that

$$P^\delta(\,\cdot\,;v^\delta;a^\delta;\Omega^\delta_\Gamma)=\frac{\omega^\delta(\,\cdot\,;\{a^\delta\};\Omega^\delta_\Gamma)}{\omega^\delta(v^\delta;\{a^\delta\};\Omega^\delta_\Gamma)}\,.$$

In the continuous setup, let Ω be some simply connected domain, $a\in P(\Omega)$ be some prime end and $v\in\operatorname{Int}\Omega$. We call $P=P(\cdot;v;a;\Omega)$ the *Poisson kernel normalized at* v, if

$$\Delta P=0\ \text{ in }\ \Omega,\qquad P\big|_{\partial\Omega\setminus\{a\}}=0,\qquad P\geqslant 0\qquad\text{and}\qquad P(v)=1$$

(note that P is uniquely defined by these conditions for arbitrary simply connected domains Ω as the conformal image of the standard Poisson kernel defined in $\mathbb{D}$).

Theorem 3.10. *The discrete Poisson kernels $P^\delta(\cdot;v^\delta;a^\delta;\Omega^\delta_\Gamma)$ are uniformly C^1–close to the continuous Poisson kernels $P(\cdot;v;a^\delta;\Omega^\delta)$ inside Ω^δ (see Definition 3.6).*



FIGURE 4. Parts of the continuous domain Ω and some discrete domain Ω^δ close to Ω in the Carathéodory topology. Marked boundary points $a^\delta \in \partial\Omega^\delta$ and $a \in P(\Omega)$ are close to each other in this topology since there are corresponding small cross-cuts near a_d. L_d denotes the closest to v arc in $\{z : |z - a_d| = d\} \cap \Omega^\delta$. The quadrilateral R_d^{3d} is shaded.

Proof. The continuous Poisson kernel is Carathéodory stable by definition, so (3.5) holds true. Thus, it is sufficient to prove pointwise convergence (3.6).

Let $(\Omega^\delta; u^\delta, v^\delta; a^\delta) \xrightarrow{\text{Cara}} (\Omega; u, v; a)$. Recall that $v^\delta \to v$ and $B(v, r) \subset \Omega^\delta$ for some $r > 0$, if δ is small enough. It follows from $P(v^\delta) = 1$ and the discrete Harnack Lemma (Proposition 2.7 (ii)) that P^δ are uniformly bounded on each compact subset of Ω. Then, due to Proposition 3.1, one can find a subsequence $\delta_k \to 0$ such that

$$P^{\delta_k}(\cdot; v^{\delta_k}; a^{\delta_k}; \Omega_\Gamma^{\delta_k}) \rightrightarrows H$$

uniformly on compact subsets of Ω, where $H \geqslant 0$ is some harmonic function in Ω. It is sufficient to prove that $H(u) = P(u; v; a; \Omega)$ for each subsequential limit H.

Let $d > 0$ be small enough. Then, there exists a crosscut $\gamma_d^a \subset B(a_d, \frac{1}{2}d)$ in Ω separating a and u, v (see Fig. 4). Moreover, one may assume that $u, v \notin \overline{B(a_d, 4d)}$ and u, v belong to the same component of $\Omega \setminus \overline{B(a_d, 4d)}$. For sufficiently small δ let

$$L_d \subset \{z : |z - a_d| = d\} \cap \Omega^\delta$$

be an arc separating v^δ and a^δ in Ω^δ (we take the arc closest to v^δ, see Fig. 4). Let Ω_d^δ denote the connected component of $\Omega^\delta \setminus L_d$ containing v^δ. Since $\Omega^\delta \xrightarrow{\text{Cara}} \Omega$ w.r.t. v and $v^\delta \to v$, one has

$$\omega(v^\delta; L_d; \Omega_d^\delta) \geqslant \text{const}(d) > 0 \tag{3.7}$$

(here and below constants const(d) do not depend on δ). Similarly, let Ω_{3d}^δ be the connected component of $\Omega^\delta \setminus L_{3d}$ containing u, v. Denote

$$M_{3d}^\delta = \max\{P^\delta(z^\delta),\ z^\delta \in \Omega_{3d}^\delta \cap \Gamma^\delta\}.$$

Since the function P^δ is discrete harmonic, one has

$$M^\delta_{3d} = P^\delta(z_0^\delta) \leqslant P^\delta(z_1^\delta) \leqslant P^\delta(z_2^\delta) \leqslant ..$$

for some nearest-neighbor path $K^\delta_{3d} = \{z_0^\delta \sim z_1^\delta \sim z_2^\delta \sim .., \ z_s^\delta \in \Omega^\delta_\Gamma\}$, starting at some $z_0^\delta \in \Omega^\delta_{3d}$. Since $P^\delta|_{\partial\Omega^\delta_\Gamma \setminus \{a^\delta\}} = 0$, the unique possibility for this path to end is a^δ.

Using (3.7), it is not hard to conclude (see Lemma 3.11 below) that the following holds true for the *continuous* harmonic measures:

$$\omega(v^\delta; K_{3d}; \Omega^\delta \setminus K_{3d}) \geqslant \omega(v^\delta; K_{3d} \cap \Omega^\delta_d; \Omega^\delta_d \setminus K_{3d}) \geqslant \mathrm{const} \cdot \omega(v^\delta; L_d; \Omega^\delta_d) \geqslant \mathrm{const}(d),$$

where K_{3d} is the corresponding polyline starting at z_0^δ and ending at a^δ. In view of Theorem 3.8, if δ is small enough, then the same inequality

$$\omega^\delta(v^\delta; K^\delta_{3d}; \Omega^\delta_\Gamma \setminus K^\delta_{3d}) \geqslant \mathrm{const}(d) > 0$$

holds true for *discrete* harmonic measures *uniformly* as $\delta \to 0$ (with smaller const(d)). Recall that $P^\delta(v^\delta) = 1$ by definition and $P^\delta(v) \geqslant M^\delta_{3d}$ along the path M^δ_{3d}. Thus,

$$M^\delta_{3d} \leqslant \mathrm{const}(d), \quad \text{if } \delta \text{ is small enough.}$$

Finally, let $z^\delta \in \Gamma^\delta \cap \Omega^\delta_{3d}$ be such that $|z^\delta - a_d| > 3d$. The weak Beurling-type estimate immediately gives

$$P^\delta(z^\delta) \leqslant \mathrm{const} \cdot \left[\frac{\mathrm{dist}(z^\delta; \partial\Omega^\delta)}{\mathrm{dist}(z^\delta; L_{3d})}\right]^\beta \cdot M^\delta_{3d} \leqslant \mathrm{const}(d) \cdot \left[\frac{\mathrm{dist}(z^\delta; \partial\Omega^\delta)}{|z^\delta - a_d| - 3d}\right]^\beta.$$

Passing to the limit as $\delta = \delta_k \to 0$, one obtains

$$H(z) \leqslant \mathrm{const}(d) \cdot \left[\frac{\mathrm{dist}(z; \partial\Omega)}{|z - a_d| - 3d}\right]^\beta \quad \text{for all } z \in \Omega_{3d} \ \ \text{suchthat} \ \ |z - a_d| > 3d.$$

Thus, $H = 0$ on $\partial\Omega \setminus \{a\}$. Since $H \geqslant 0$ and $H(v) = 1$, it gives $H = P(\cdot; v; a; \Omega)$. ☐

Lemma 3.11. *Let $\Omega \subset \mathbb{C}$ be some simply connected domain, $v \in \mathrm{Int}\,\Omega$ and $a \in P(\Omega)$. Let $L_d \subset \{z : |z - a_d| = d\} \cap \Omega$ be the arc separating v and a that is closest to v, and Ω_d be the connected component of $\Omega \setminus L_d$ containing v. Let K_{3d} be some path connecting L_{3d} and L_d inside the conformal quadrilateral $R^{3d}_d = \Omega_d \setminus \overline{\Omega_{3d}}$ (see Fig. 4). Then*

$$\omega(v; K_{3d}; \Omega_d \setminus K_{3d}) \geqslant \mathrm{const} \cdot \omega(v; L_d; \Omega_d)$$

for some absolute positive constant.

Proof. Note that $\omega(v; L_d; \Omega_d) \leqslant \omega(v; L_d; \Omega_d \setminus K_{3d}) + \omega(v; K_{3d}; \Omega_d \setminus K_{3d})$. Thus, it is sufficient to prove that

$$\omega(v; L_d; \Omega_d \setminus K_{3d}) \leqslant \mathrm{const} \cdot \omega(v; K_{3d}; \Omega_d \setminus K_{3d}).$$

Furthermore, monotonicity arguments give

$$\omega(v; L_d; \Omega_d \setminus K_{3d}) \leqslant \int_{L_{2d}} \omega(z; L_d \cup L_{3d}; R^{3d}_d \setminus K_{3d}) \cdot \omega(v; |dz|; \Omega_{2d} \setminus K_{3d})$$

and, in a similar manner,

$$\omega(v; K_{3d}; \Omega_d \setminus K_{3d}) \geqslant \int_{L_{2d}} \omega(z; K_{3d}; R^{3d}_d \setminus K_{3d}) \cdot \omega(v; |dz|; \Omega_{2d} \setminus K_{3d}).$$

Let $L_d = A_d B_d$ and so on (see Fig. 4). Monotonicity arguments give

$$\omega(z; K_{3d}; R_d^{3d} \setminus K_{3d}) \geqslant \min\left\{\omega(z; A_{3d}A_d; R_d^{3d}),\, \omega(z; B_dB_{3d}; R_d^{3d})\right\}.$$

Thus, it is sufficient to prove that

$$\omega(z; A_dB_d \cup B_{3d}A_{3d}; R_d^{3d}) \leqslant \mathrm{const} \cdot \min\left\{\omega(z; A_{3d}A_d; R_d^{3d}),\, \omega(z; B_dB_{3d}; R_d^{3d})\right\}$$

for all $z \in L_{2d}$. Due to the conformal invariance of harmonic measure, the last estimate follows from the uniform bounds on the extremal distances (conformal modulii of quadrilaterals)

$$\lambda_{A_{nd}B_{nd}B_{md}A_{md}}(A_{nd}A_{md}\,;\, B_{md}B_{nd}) \geqslant \frac{1}{2\pi}\log\frac{m}{n}, \qquad 1 \leqslant n < m \leqslant 3. \qquad \square$$

3.5. **Convergence for the Poisson kernel normalized at a boundary point.** Recall that $\mathbb{H}^\delta$ denotes the discretization of the upper half-plane $\mathbb{H} = \{z : \operatorname{Im} z > 0\}$ (i.e., the set of all faces, edges and vertices that intersect $\mathbb{H}$, see Fig. 2B). As for bounded domains, denote by $\omega^\delta(u^\delta; \{x^\delta\}; \mathbb{H}^\delta_\Gamma)$ the probability of the event that the random walk starting at $u^\delta \in \mathbb{H}^\delta_\Gamma$ first hits the boundary $\partial\mathbb{H}^\delta_\Gamma$ at a vertex $x^\delta \in \partial\mathbb{H}^\delta_\Gamma$. It is easy to see (e.g., using the unboundedness of the free Green's function (2.3) or Proposition 2.10) that

$$\sum_{x^\delta \in \partial\mathbb{H}^\delta_\Gamma} \omega^\delta(u^\delta; \{x^\delta\}; \mathbb{H}^\delta_\Gamma) = 1.$$

Let

$$\Im^\delta(u^\delta) = \operatorname{Im} u^\delta - \sum_{x^\delta \in \partial\mathbb{H}^\delta_\Gamma} \operatorname{Im} x^\delta \cdot \omega^\delta(u^\delta; \{x^\delta\}; \mathbb{C}^\delta_+) \quad \text{for} \quad u^\delta \in \mathbb{H}^\delta_\Gamma. \qquad (3.8)$$

The function $\Im^\delta$ is discrete harmonic in $\mathbb{H}^\delta_\Gamma$, $\Im^\delta = 0$ on $\partial\mathbb{H}^\delta_\Gamma$ and $|\Im^\delta(u^\delta) - \operatorname{Im} u^\delta| \leqslant 2\delta$ for all $u^\delta \in \mathbb{H}^\delta_\Gamma$ (note that these conditions define $\Im^\delta$ uniquely). In particular, if $\operatorname{Im} u^\delta \in [3\delta, 5\delta]$, then $\Im^\delta(u^\delta) \asymp \delta$. Since $\Im^\delta$ is discrete harmonic and nonnegative, it implies

$$\Im^\delta(x^\delta_{\mathrm{int}}) \asymp \delta \qquad \text{for all} \quad x^\delta = (x^\delta; x^\delta_{\mathrm{int}}) \in \partial\mathbb{H}^\delta_\Gamma.$$

Below we need some simple technical estimates for the discrete harmonic measure in the rectangle at a point lying near the boundary. Let $o^\delta \in \partial\mathbb{H}^\delta_\Gamma$ be the boundary vertex closest to 0,

$$R(S,T) = (-S; S) \times (0; T) \subset \mathbb{C}, \qquad S, T > 0,$$

$R^\delta_\Gamma(S,T) \subset \Gamma$ denote the discretization of this open rectangle, and

$$\begin{aligned} L^\delta_\Gamma(S) \quad &= \quad \left\{v^\delta \in \partial R^\delta_\Gamma(S,T) : \operatorname{Im} v^\delta \leqslant 0\right\}, \\ U^\delta_\Gamma(S,T) \quad &= \quad \left\{v^\delta \in \partial R^\delta_\Gamma(S,T) : \operatorname{Im} v^\delta \geqslant T\right\}, \\ V^\delta_\Gamma(S,T) \quad &= \quad \left\{v^\delta \in \partial R^\delta_\Gamma(S,T) : |\operatorname{Re} v^\delta| \geqslant S\right\} \end{aligned}$$

be the lower, upper and vertical parts of the boundary $\partial R^\delta_\Gamma(S,T)$ (see Fig. 2B).

Lemma 3.12. *Let $s \geqslant 2t > 0$. If δ is small enough, then*

$$\omega^\delta(o^\delta_{\mathrm{int}}; U^\delta_\Gamma(s,t); R^\delta_\Gamma(s,t)) \asymp \delta/t \quad \textit{and} \quad \omega^\delta(o^\delta_{\mathrm{int}}; V^\delta_\Gamma(s,t); R^\delta_\Gamma(s,t)) \leqslant \mathrm{const} \cdot \delta t/s^2.$$

Proof. We defer the proof until after that of Theorem 3.13. $\square$

From now on, we consider only discrete domains $(\Omega^\delta_\Gamma; a^\delta)$, $a^\delta \in \partial\Omega^\delta_\Gamma$ such that

$$R^\delta_\Gamma(S,T) \subset \Omega^\delta_\Gamma, \qquad L^\delta_\Gamma(S) \subset \partial\Omega^\delta_\Gamma \qquad \text{and} \qquad a^\delta \notin L^\delta_\Gamma(S) \tag{3.9}$$

for some $S,T>0$. All continuous domains $(\Omega; a)$ appearing as Carathéodory limits of these $(\Omega^\delta; a^\delta)$ satisfy

$$R(S,T) \subset \Omega, \qquad [-\tfrac{1}{2}S; \tfrac{1}{2}S] \subset \partial\Omega \qquad \text{and} \qquad a \notin [-\tfrac{1}{2}S; \tfrac{1}{2}S]. \tag{3.10}$$

For continuous domains $(\Omega; a)$ satisfying (3.10), we define the *Poisson kernel* $P_0(\cdot) = P_0(\cdot; a; \Omega)$ *normalized at* 0 as the (unique) solution of the following problem:

$$\Delta P_0 = 0 \ \text{ in } \ \Omega, \qquad P_0\big|_{\partial\Omega\setminus\{a\}} = 0, \qquad P_0 \geqslant 0 \qquad \text{and} \qquad \partial_y P_0(0) = 1.$$

For discrete domains $(\Omega^\delta_\Gamma; a^\delta)$ satisfying (3.9), we call $P^\delta_{o^\delta} = P^\delta_{o^\delta}(\cdot; a^\delta; \Omega^\delta_\Gamma)$ the *discrete Poisson kernel normalized at* o^δ, if

$$\Delta^\delta P^\delta_{o^\delta} = 0 \ \text{ in } \ \Omega^\delta_\Gamma, \qquad P^\delta_{o^\delta}\big|_{\partial\Omega^\delta_\Gamma\setminus\{a\}} = 0 \qquad \text{and} \qquad P^\delta_{o^\delta}(o^\delta_{\mathrm{int}}) = \Im(o^\delta_{\mathrm{int}}).$$

Note that

$$P^\delta_{o^\delta}(\cdot; a^\delta; \Omega^\delta_\Gamma) = \omega^\delta(\cdot; a^\delta; \Omega^\delta_\Gamma) \cdot \frac{\Im^\delta(o^\delta_{\mathrm{int}})}{\omega^\delta(o^\delta_{\mathrm{int}}; a^\delta; \Omega^\delta_\Gamma)}\,.$$

Theorem 3.13. *The discrete Poisson kernels $P^\delta_{o^\delta}(\cdot; a^\delta; \Omega^\delta_\Gamma)$ defined for the class of discrete domains satisfying (3.9) with some $S,T>0$ are uniformly C^1–close in $\widetilde{\Omega}^\delta$ (see Definition 3.6) to the continuous Poisson kernels $P_0(\cdot; a^\delta; \widetilde{\Omega}^\delta)$, where $\widetilde{\Omega}^\delta$ denotes the modified polygonal domain Ω^δ with the part of the boundary $L^\delta(\frac{1}{2}S) \subset \partial\Omega^\delta$ replaced by the segment $[-\frac{1}{2}S, \frac{1}{2}S]$. The rate of uniform convergence depends on S,T.*

Proof. The continuous Poisson kernel P_0 is Carathéodory stable, so (3.5) holds true. Thus, it is sufficient to check (3.6). Without loss of generality, we assume that $S \geqslant 2T$.

Let $(\Omega^\delta; u^\delta; a^\delta) \xrightarrow{\text{Cara}} (\Omega; u; a)$. Denote by $T^\delta \in U^\delta_\Gamma(\frac{1}{2}S, \frac{1}{2}T)$ the vertex closest to the point $\frac{i}{2}T \in \mathbb{C}$. Due to the discrete Harnack Lemma (Proposition 2.7 (ii)), the values of $P^\delta_{o^\delta}$ on $U^\delta_\Gamma(\frac{1}{2}S, \frac{1}{2}T)$ are uniformly comparable with $P^\delta_{o^\delta}(T^\delta)$. Using Lemma 3.12, one obtains

$$\delta \asymp P^\delta_{o^\delta}(o^\delta_{\mathrm{int}}) \geqslant \mathrm{const} \cdot P^\delta_{o^\delta}(T^\delta) \cdot \omega^\delta(o^\delta_{\mathrm{int}}; U^\delta_\Gamma(\tfrac{1}{2}S, \tfrac{1}{2}T); R^\delta_\Gamma(\tfrac{1}{2}S, \tfrac{1}{2}T)) \geqslant \mathrm{const} \cdot \frac{\delta}{T} \cdot P^\delta_{o^\delta}(T^\delta)$$

and so the values $P^\delta_{o^\delta}(T^\delta)$ are uniformly bounded. Hence, applying the discrete Harnack Lemma once more, one derives the uniform boundedness of $P^\delta_{o^\delta}(\cdot; a^\delta; \Omega^\delta_\Gamma)$ on each compact subset of Ω. Then, due to Proposition 3.1, one can find a subsequence $\delta_k \to 0$ such that

$$P^{\delta_k}_{o^\delta}(\cdot; a^{\delta_k}; \Omega^{\delta_k}_\Gamma) \rightrightarrows H$$

uniformly on compact subsets of Ω, where $H \geqslant 0$ is some harmonic in Ω function. We need to prove that $H(u) = P_0(u; a; \Omega)$ for each subsequential limit H.

Repeating the arguments given in the proof of Theorem 3.10, one obtains that, first, for each $r>0$ the functions $P^\delta_{o^\delta}$ are uniformly bounded *everywhere* in $\Omega^\delta_r \cap \Gamma^\delta$ (in particular, everywhere in the rectangle $R^\delta_\Gamma(S,T)$) and, second, $H = 0$ on $\partial\Omega \setminus \{a\}$. Therefore, due to $H \geqslant 0$,

$$H = \mu P_0(\cdot; a; \Omega) \quad \text{for some} \ \ \mu \geqslant 0.$$

Let

$$Q^\delta(\cdot) = P^\delta_{o^\delta}(\cdot; a^\delta; \Omega^\delta_\Gamma) - \Im^\delta(\cdot).$$

By definition, the function Q^δ is discrete harmonic in $R^\delta_\Gamma(S,T)$, $Q^\delta = 0$ on the lower boundary $L^\delta_\Gamma(S)$, $Q^\delta(o^\delta_{\mathrm{int}}) = 0$ and

$$Q^\delta(v) \rightrightarrows \mu P_0(v; a; \Omega) - \operatorname{Im} v \qquad \text{as } \ \delta = \delta_k \to 0$$

uniformly on compact subsets of $R(S,T)$.

Since $P_0|_{[-S,S]} = 0$ and $(P_0)'_y(0) = 1$, one has $P_0(x+iy) = y + O(xy + y^2)$ as $x, y \to 0$. Thus, for any fixed $s < S$ and $t < T$, $2t \leqslant s$, the following hold true:

$$\begin{aligned} &Q^\delta = 0 \quad \text{on} \ \ L^\delta_\Gamma(s), \qquad\qquad |Q^\delta| \leqslant \text{const} \quad \text{on} \ \ V^\delta_\Gamma(s,t), \\ &Q^\delta \rightrightarrows (\mu-1)t + O(st) \ \ \text{as } \delta \to 0 \ \ \text{uniformly on} \ \ U^\delta_\Gamma(t,s). \end{aligned}$$

The normalization $Q^\delta(o^\delta_{\mathrm{int}}) = 0$ and Lemma 3.12 give

$$|(\mu-1)t + O(st) + o(1)| \cdot \delta/t \leqslant \text{const} \cdot \delta t/s^2$$

as $\delta \to 0$. So, for any fixed s and t, one has $|\mu-1| \leqslant \text{const} \cdot (s + t/s^2)$. Thus, $\mu = 1$. ☐

Proof of Lemma 3.12. We consider two quadratic harmonic polynomials

$$h_1(x+iy) = \left(1 - \frac{(t+2\delta)^2}{s^2}\right) \cdot \frac{y}{t+2\delta} + \frac{y^2 - x^2}{s^2}$$

and

$$h_2(x+iy) = \left(1 + \frac{(t+2\delta)^2}{s^2}\right) \cdot \frac{y+2\delta}{t+2\delta} + \frac{x^2 - (y+2\delta)^2}{s^2}.$$

Their restrictions on Γ are discrete harmonic due to Lemma 2.2 (i). Moreover, it is easy to check that

$$\begin{aligned} &h_1(v^\delta) \leqslant 0 \leqslant h_2(v^\delta), \quad \text{if} \quad v^\delta \in L^\delta_\Gamma(s) \cup V^\delta_\Gamma(s,t), \\ &h_1(v^\delta) \leqslant 1 \leqslant h_2(v^\delta), \quad \text{if} \quad v^\delta \in U^\delta_\Gamma(s,t). \end{aligned}$$

Hence, $h_1(v^\delta) \leqslant \omega^\delta(v^\delta; U^\delta_\Gamma(s,t); R^\delta_\Gamma(s,t)) \leqslant h_2(v^\delta)$ for all $v^\delta \in R^\delta_\Gamma(s,t)$. Thus,

$$\omega^\delta(v^\delta; U^\delta_\Gamma(s,t); R^\delta_\Gamma(s,t)) \asymp \delta/t,$$

if, say, $\operatorname{Im} v^\delta \in [2\delta, 4\delta]$ and $|\operatorname{Re} v^\delta| \leqslant 4\delta$. Since ω^δ is discrete harmonic, it implies

$$\omega^\delta(o^\delta_{\mathrm{int}}; U^\delta_\Gamma(s,t); R^\delta_\Gamma(s,t)) \asymp \delta/t.$$

The upper bound for $\omega^\delta(v^\delta; V^\delta_\Gamma(s,t); R^\delta_\Gamma(s,t))$ follows by the similar consideration of the quadratic harmonic polynomial

$$h_3(x+iy) = \frac{x^2 + (y+2\delta)(t+2\delta-y)}{s^2}$$

which is nonnegative on $L^\delta_\Gamma(s) \cup U^\delta_\Gamma(s,t)$ and no less than 1 on $V^\delta_\Gamma(s,t)$. ☐

A. Appendix

A.1. **Kenyon's asymptotics for the Green's function and the Cauchy kernel.** Below we give a sketch of R. Kenyon's [Ken02] arguments with some small improvement concerning the estimates of the integrands for intermediate λ. See also [Bück08].

Proof of the Theorem 2.5. Following J. Ferrand [Fer44] and Ch. Mercat [Mer07] we introduce *discrete exponentials*

$$e(\lambda,u;u_0)=\prod_{j=0}^{2k-1}\frac{1+\frac{\lambda}{2}(u_k-u_{k-1})}{1-\frac{\lambda}{2}(u_k-u_{k-1})}, \tag{A.1}$$

where $P_{u_0u}=u_0u_1u_2...u_{2k-1}u_{2k}$ is a path from u_0 to $u_{2k}=u$ on the corresponding rhombic embedding (so $u_{2j}\in\Gamma$ and $u_{2j-1}\in\Gamma^*$ for all j). It's easy to see that this definition does not depend on the choice of the path. Since the angles of rhombi are bounded from 0 and π, one can choose P_{u_0u} so that the following condition holds:

> for all j either (a) $|\arg(u_{j+1}-u_j)-\arg(u-u_0)|<\frac{\pi}{2}$ or (b) u_j and u_{j+2} are opposite vertices of some rhombus and $|\arg(u_{j+2}-u_j)-\arg(u-u_0)|<\frac{\pi}{2}$.
> In particular, $\arg(u_{j+1}-u_j)-\arg(u-u_0)\in(-\pi,\pi)$ for all j.

Define (see R. Kenyon [Ken02] and A. Bobenko, Ch. Mercat and Yu. Suris [BMS05])

$$\widetilde{G}_\Gamma(u;u_0)=\frac{1}{8\pi^2 i}\int_C\frac{\log\lambda}{\lambda}\,e(\lambda,u;u_0)\,d\lambda. \tag{A.2}$$

where C is a curve which runs counter clockwise around the disc of (large) radius R from the angle $\arg\overline{(u-u_0)}-\pi$ to $\arg\overline{(u-u_0)}+\pi$, then along the segment $e^{i\arg\overline{(u-u_0)}}[-R,-r]$, then clockwise around the disc of (small) radius r and then back along the same segment (the integral does not depend on the log branch, since $e(0,u;u_0)=e(\infty,u;u_0)=1$).

This function is discrete harmonic away from u_0 since all discrete exponentials are harmonic (as functions of u) and one can use the same contour of integration for all $u_s\sim u$. Furthermore, $\widetilde{G}(u_0;u_0)=0$ and, by straightforward computation,

$$\widetilde{G}_\Gamma(u_s;u_0)=\frac{\theta_s\cot\theta_s}{\pi},\quad\text{if}\quad u_s\sim u,\qquad\text{so}\qquad \mu_\Gamma^\delta(u_0)\cdot\Delta^\delta\widetilde{G}_\Gamma(u_0;u_0)=1.$$

Rotating and scaling the plane, one may assume that $\arg(u-u_0)=0$ and $\delta=1$. It's easy to see that the contribution of intermediate $\lambda=-t<0$ to the integral in (A.2) is exponentially small. Indeed, in case (a) one has

$$\left|\frac{1-\frac{t}{2}e^{i\beta_j}}{1+\frac{t}{2}e^{i\beta_j}}\right|^2=1-\frac{8t\cos\beta_j}{t^2+4t\cos\beta_j+4}\leqslant\exp\left[-\frac{8t\cos\beta_j}{(t+2)^2}\right],$$

where $\beta_j=\arg(u_{j+1}-u_j)$ and $\cos\beta_j=\operatorname{Re}(u_{j+1}-u_j)>0$. Similarly, in case (b),

$$\left|\frac{(1-\frac{t}{2}e^{i\beta_j})(1-\frac{t}{2}e^{i\beta_{j+1}})}{(1+\frac{t}{2}e^{i\beta_j})(1+\frac{t}{2}e^{i\beta_{j+1}})}\right|^2\leqslant\exp\left[-\frac{8t(\cos\beta_j+\cos\beta_{j+1})}{(t+2)^2}\right],$$

due to $\cos\beta_j+\cos\beta_{j+1}=\operatorname{Re}(u_{j+2}-u_j)>0$. Thus,

$$|e(-t,u;u_0)|\leqslant\exp\left[-\frac{4t(u-u_0)}{(t+2)^2}\right]$$

and the asymptotics of (A.2) as $|u-u_0|\to\infty$ are determined by the asymptotics of $e(\lambda,u;u_0)$ near 0 and ∞. Some version of the Laplace method (see [Ken02] and [Bück08]) gives

$$\widetilde{G}_\Gamma(u;u_0)=\frac{1}{2\pi}\log|u-u_0|+\frac{\gamma_{\mathrm{Euler}}+\log 2}{2\pi}+O(|u-u_0|^{-2}),$$

where the remainder is of order $|u-u_0|^{-2}$ due to

$$\frac{d^2}{d\lambda^2}\log(e(\lambda,u;u_0))\Big|_{\lambda=0}=\frac{d^2}{d\lambda^2}\log(e(\lambda,u;u_0))\Big|_{\lambda=\infty}=0.$$

The uniqueness of $G(\cdot;u_0)$ (and $G=\operatorname{Re}G$) easily follows by the Harnack inequality (Corollary 2.8). Indeed, the difference $G=G_1(\cdot;u_0)-G_2(\cdot;u_0)$ is discrete harmonic on Γ and $\max_{|u|\leqslant R}|H(u)|/R\to 0$ as $u\to\infty$, so $G(u)=G(u_0)=0$ for all u. □

Proof of Theorem 2.14. As in Theorem 2.5, $K(\cdot;z_0)$ can be constructed using (modified) discrete exponentials. Similarly to (A.1) denote

$$e(\lambda,v_j^0;z_0)=\frac{1}{(1-\frac{\lambda}{2}(v_j^0-v_{j-1}^0))(1-\frac{\lambda}{2}(v_j^0-v_{j+1}^0))},\quad j=1,2,3,4, \tag{A.3}$$

for the vertices of the rhombus $v_1^0v_2^0v_3^0v_4^0$ centered at z_0 and, by induction,

$$\frac{e(\lambda,w;z_0)}{e(\lambda,u;z_0)}=\frac{1+\frac{\lambda}{2}(w-u)}{1-\frac{\lambda}{2}(w-u)}\quad\text{for all}\ \ w\sim u,\ u\in\Gamma,\ w\in\Gamma^*.$$

As above, all $e(\lambda,\cdot;z_0)$ are discrete holomorphic on Λ. Define (see [Ken02])

$$K(v;z_0)=\frac{1}{\pi}\int_{-\overline{(v-z_0)}\infty}^{0}e(\lambda,v;z_0)d\lambda,$$

where the integral being, say, along the ray $\arg\zeta=\arg\overline{(v-z_0)}\pm\pi$ (taking the path from z_0 to u as in the proof of Theorem 2.5, one guarantees that all poles of $e(\cdot,v;z_0)$ are in $|\arg\lambda-\arg(v-z_0)|<\pi$). Then $K(\cdot;z_0)$ is holomorphic everywhere except z_0. The straightforward calculations give

$$\mu_{z_0v_j^0}\cdot K(v_j^0;z_0)=\frac{\arg(v_j^0-v_{j-1}^0)-\arg(v_j^0-v_{j+1}^0)}{\pi},\quad\text{so}\quad \mu_\diamondsuit^\delta(z_0)\cdot[\overline{\partial}^\delta K](z_0;z_0)=1.$$

Let $\delta=1$. As in Theorem 2.5, the integrand is exponentially small for intermediate λ. One has

$$e(\lambda,v;z_0)=\exp\left[\lambda(v-z_0)+O(|\lambda|^2|v-z_0|)\right],\qquad \lambda\to 0,$$

and

$$e(\lambda,v;z_0)=\frac{4\overline{\tau}^2}{\lambda^2}\cdot\exp\left[\frac{4\overline{(v-z_0)}}{\lambda}+O\left(\frac{|v-z_0|}{|\lambda|^2}\right)\right],\quad \lambda\to\infty,$$

where $\tau=e^{i\arg(u_0^{(1)}-u_0^{(2)})}$, if $v\in\Gamma$, and $\tau=e^{i\arg(w_0^{(1)}-w_0^{(2)})}$, if $v\in\Gamma^*$ (the factor $\overline{\tau}^2$ comes from (A.3)). Summarizing, one arrives at

$$K(u;z_0)=\frac{1}{\pi}\left[\frac{1}{v-z_0}+\frac{\overline{\tau}^2}{\overline{v}-\overline{z}_0}\right]+O\left(\frac{1}{|v-z_0|^2}\right)=\frac{2}{\pi}\Pr\left[\frac{1}{v-z_0};\overline{\tau}\right]+O\left(\frac{1}{|v-z_0|^2}\right).$$

Finally, $K(\cdot,z_0)$ is unique due to Corollary 2.8. □

A.2. **Proof of the discrete Harnack Lemma.** Below we recall the modification of R. Duffin's arguments [Duf53] given by U. Bücking [Bück08]. For the next it is important that the remainder in (2.3) is of order $\delta^2|u-u_0|^{-2}$ (and not just $\delta|u-u_0|^{-1}$).

Proposition A.1. *Let* $u_0\in\Gamma$ *and* $R\geqslant\delta$. *Then*

$$\omega^\delta(u_0;\{a\};B_\Gamma^\delta(u_0,R))\asymp\frac{\delta}{R}\qquad\textit{for all } a\in\partial B_\Gamma^\delta(u_0,R),$$

i.e., $\mathrm{const}_1\cdot\delta/R\leqslant\omega^\delta(u_0;\{a\};\Omega_\Gamma^\delta)\leqslant\mathrm{const}_2\cdot\delta/R$ *for some positive absolute constants.*

Proof. One has $R\leqslant|a-u_0|\leqslant R+2\delta$ for all $a\in\partial B_\Gamma^\delta(u_0,R)$. Therefore, (2.3) gives

$$\left|G_{B_\Gamma^\delta(u_0,R)}(u;u_0)-\frac{1}{2\pi}\log\frac{|u-u_0|}{R}\right|\leqslant\frac{\delta}{\pi R}+\mathrm{const}\cdot\left(\frac{\delta^2}{|u-u_0|^2}+\frac{\delta^2}{R^2}\right)$$

for all $u\neq u_0$. In particular, if R/δ is large enough, then

$$|G_{B_\Gamma^\delta(u_0,R)}(u;u_0)|\asymp\frac{\delta}{R}\quad\text{for all } u\in B_\Gamma^\delta(u_0,R):\ R-5\delta\leqslant|u-u_0|\leqslant R-3\delta.$$

Since Green's function is discrete harmonic and nonpositive near $\partial B_\Gamma^\delta(u_0,R)$, the same holds true for all $a_{\mathrm{int}}:(a;a_{\mathrm{int}})\in\partial B_\Gamma^\delta(u_0,R)$. In view of (2.5), this gives the result for sufficiently large R/δ. For small (i.e. comparable to δ) radii R the claim is trivial, since the random walk can reach a starting from u_0 in a finite number of steps. □

Proposition A.2 (mean value property). *Let* $H:B_\Gamma^\delta(u_0,R)\to\mathbb{R}$ *be nonnegative discrete harmonic function. Then*

$$\left|H(u_0)-\frac{1}{\pi R^2}\sum_{u\in\mathrm{Int}\,B_\Gamma^\delta(u_0,R)}H(u)\mu_\Gamma^\delta(u)\right|\leqslant\mathrm{const}\cdot\frac{\delta H(u_0)}{R}.$$

Proof. Let

$$F(u)=G_\Gamma(u;u_0)-\frac{\log R}{2\pi}+\frac{R^2-|u-u_0|^2}{4\pi R^2},\quad u\in B_\Gamma^\delta=B_\Gamma^\delta(u_0,R).$$

Note that $[\Delta^\delta F](u)=-(\pi R^2)^{-1}$ for all $u\neq u_0$ (see Lemma 2.2(i)). Using (2.3), it is easy to see that $F(u)=O(\delta^2/R^2)$, if $|u-R|\leqslant\mathrm{const}\cdot\delta$. The discrete Green's formula gives

$$H(u_0)-\frac{1}{\pi R^2}\sum_{u\in\mathrm{Int}\,B_\Gamma^\delta}H(u)\mu_\Gamma^\delta(u)\ =\sum_{a\in\partial B_\Gamma^\delta}\tan\theta_{aa_{\mathrm{int}}}\cdot[F^\pm(a)H(a_{\mathrm{int}})-F^\pm(a_{\mathrm{int}})H(a)],$$

where the functions $F^\pm=F\pm\mathrm{const}\cdot\delta^2/R^2$ are positive/negative, respectively, near $\partial B_\Gamma^\delta(u_0,R)$. Using $H\geqslant0$, one obtains

$$-\mathrm{const}\cdot\frac{\delta^2}{R^2}\sum_{a\in\partial B_\Gamma^\delta}H(a)\leqslant H(u_0)-\frac{1}{\pi R^2}\sum_{u\in\mathrm{Int}\,B_\Gamma^\delta}H(u)\mu_\Gamma^\delta(u)\leqslant\mathrm{const}\cdot\frac{\delta^2}{R^2}\sum_{a\in\partial B_\Gamma^\delta}H(a_{\mathrm{int}}),$$

Both sums are comparable to $\delta H(u_0)/R$ due to Proposition A.1. □

Proof of Proposition 2.7. (i) Applying the mean value property for $B^\delta_\Gamma(u_0,R)$ and $B^\delta_\Gamma(u_1,R{-}2\delta)$ and taking into account $H(u_1)\leqslant \mathrm{const}\cdot H(u_0)$, one obtains

$$\pi R^2H(u_0)-\pi(R{-}2\delta)^2H(u_1) \quad = \sum_{u\in B^\delta_\Gamma(u_0,R)\setminus B^\delta_\Gamma(u_1,R-2\delta)} H(u)\mu^\delta_\Gamma(u)+O(\delta RH(u_0)).$$

Proposition A.1 gives

$$\sum_{u\in B^\delta_\Gamma(u_0,R)\setminus B^\delta_\Gamma(u_1,R-2\delta)} H(u)\mu^\delta_\Gamma(u) \quad\asymp \sum_{u\in B^\delta_\Gamma(u_0,R)\setminus B^\delta_\Gamma(u_1,R-2\delta)} \delta^2H(u) \;=\; O(\delta RH(u_0)),$$

so $R^2(H(u_1){-}H(u_0))=O(\delta RH(u_0))$.

(ii) Let $u_1=v_0v_1v_2...v_{k-1}v_k=u_2$ be some path connecting u_1 and u_2 inside $B^\delta_\Gamma(u_0,r)$ (one can choose a path such that $k\leqslant \mathrm{const}\cdot r/\delta$). Since $B^\delta_\Gamma(v_j,R{-}r)\subset B^\delta_\Gamma(u_0,R)$,

$$\frac{H(u_2)}{H(u_1)}=\prod_{j=0}^{k-1}\frac{H(v_{j+1})}{H(v_j)}\leqslant\left[1+\mathrm{const}\cdot\frac{\delta}{R-r}\right]^{\mathrm{const}\cdot r/\delta}\leqslant\;\exp\left[\mathrm{const}\cdot\frac{r}{R-r}\right]. \qquad \square$$

References

[BMS05] A. Bobenko, Ch. Mercat, Yu. Suris. Linear and nonlinear theories of discrete analytic functions. Integrable structure and isomonodromic Green's function. *J. Reine Angew. Math.*, 583:117–161, 2005.

[Bück08] U. Bücking. Approximation of conformal mappings by circle patterns. `arXiv:0806.3833`

[ChSm08] D. Chelkak, S. Smirnov. Universality and conformal invariance in the Ising model. In Oberwolfach report No. 25/2008 (Stochastic Analysis Workshop).

[CFL28] R. Courant, K. Friedrichs, H. Lewy. Über die partiellen Differenzengleichungen der mathematischen Physik. *Math. Ann.*, 100:32–74, 1928.

[Duf53] R. J. Duffin. Discrete potential theory. *Duke Math. J.*, 20:233–251, 1953.

[Duf68] R. J. Duffin. Potential theory on a rhombic lattice. *J. Combinatorial Theory*, 5:258–272, 1968.

[DN03] I. A. Dynnikov, S. P. Novikov, Geometry of the triangle equation on two-manifolds, *Mosc. Math. J.*, 3(2):419-438, 2003.

[Fer44] J. Ferrand, Fonctions préharmoniques et fonctions préholomorphes. *Bull. Sci. Math.*, 2nd ser., 68:152–180, 1944.

[L-F55] J. Lelong-Ferrand. *Représentation conforme et transformations à intégrale de Dirichlet bornée.* Gauthier-Villars, Paris, 1955.

[Ken02] R. Kenyon. The Laplacian and Dirac operators on critical planar graphs. *Invent. Math.*, 150:409–439, 2002.

[KSch04] R. Kenyon, J.-M. Schlenker. Rhombic embeddings of planar quad-graphs, *Trans. AMS*, 357:3443–3458, 2004.

[LSchW04] G. F. Lawler, O. Schramm, W. Werner. Conformal invariance of planar loop-erased random walks and uniform spanning trees. *Ann. Probab.*, 32(1B):939–995, 2004.

[Mer01] Ch. Mercat. Discrete Riemann Surfaces and the Ising Model. *Comm. Math. Phys.*, 218(1):177–216, 2001.

[Mer02] Ch. Mercat. Discrete Polynomials and Discrete Holomorphic Approximation. `arXiv: math-ph/0206041`

[Mer07] Ch. Mercat. Discrete Riemann Surfaces. Handbook of Teichmüller theory. Vol. I, 541–575. Zürich: European Mathematical Society (EMS), 2007.

[Pom92] Ch. Pommerenke. *Boundary Behaviour of Conformal Maps*, A Series of Comprehensive Studies in Mathematics 299. Springer-Verlag, Berlin, 1992.

[Sm06] S. Smirnov. Towards conformal invariance of 2D lattice models. Proceedings of the international congress of mathematicians (ICM), Madrid, Spain, August 22–30, 2006. Vol. II: Invited lectures, 1421-1451. Zürich: European Mathematical Society (EMS), 2006.

[Sm07] S. Smirnov. Conformal invariance in random cluster models. I. Holomorphic fermions in the Ising model. `arXiv:0708.0039` *Annals Math.*, to appear.